

On ω -categorical groups and rings with NIP

Krzysztof Krupiński*

Abstract

We show that ω -categorical rings with NIP are nilpotent-by-finite. We prove that an ω -categorical group with NIP and fsg is nilpotent-by-finite. We also notice that an ω -categorical group with at least one strongly regular type is abelian. Moreover, we get that each ω -categorical, characteristically simple p -group with NIP has an infinite, definable abelian subgroup. Assuming additionally the existence of a non-algebraic, generically stable over \emptyset type, such a group is abelian.

0 Introduction

Recall that a first order structure M in a countable language is said to be ω -categorical if, up to isomorphism, $Th(M)$ has at most one model of cardinality \aleph_0 . There is a long history of results describing the structure of ω -categorical groups and rings. However, many questions in this area are still wide open. It follows easily that each countable, ω -categorical group has a finite series of characteristic subgroups in which all successive quotients are characteristically simple groups. On the other hand, Wilson (see [19, 1]) proved

Fact 0.1 *For each countably infinite, ω -categorical, characteristically simple group H , one of the following holds.*

1. H is an elementary abelian p -group for some prime p .
2. $H \cong B(F)$ or $H \cong B^{-1}(F)$ for some non-abelian, finite, simple group F , where $B(F)$ is the group of all continuous functions from the Cantor set \mathcal{C} to F , and $B^{-1}(F)$ is the subgroup of $B(F)$ consisting of the functions f such that $f(x_0) = e$ for a fixed element $x_0 \in \mathcal{C}$.
3. H is a perfect p -group.

*Research supported by the Polish Government grant N N201 545938

⁰2000 Mathematics Subject Classification: 03C45, 03C35, 20A15

⁰Key words and phrases: ω -categorical group, ω -categorical ring, non independence property

Moreover, it is conjectured that (iii) is not realized.

As to ω -categorical rings in general, we know that their Jacobson radical is nilpotent (see [3, Lemma 1.3] and [4]). However, there are examples of infinite, ω -categorical rings which are semisimple (i.e. with trivial Jacobson radical) and so not nilpotent-by-finite [3].

Interesting questions arise when one imposes additional model-theoretic restrictions (e.g. stability or simplicity) on our ω -categorical group or ring. In the superstable [or, more generally, supersimple] ω -categorical context, everything is clear: groups are [(finite central)-by-]abelian-by-finite [17, 6]; rigs are [(finite null)-by-]null-by-finite [12]. In the stable [or, more generally, NSOP] situation, we only know that ω -categorical groups are nilpotent-by-finite [14], and ω -categorical rings are nilpotent-by-finite [3, 11], too. It is an open question whether ω -categorical stable groups are abelian-by-finite and whether ω -categorical stable rings are null-by-finite.

Our motivating problem is to describe the structure of ω -categorical groups and rings satisfying NIP. Reasonable conjectures seems to be:

Conjecture 0.2 *Each ω -categorical group with NIP is nilpotent-by-finite.*

Conjecture 0.3 *Each ω -categorical ring with NIP is nilpotent-by-finite.*

In this paper, we prove Conjecture 0.3. As to Conjecture 0.2, we prove it under the additional assumption that the group has fsg (finitely satisfiable generics).

One of the ingredients of our proof of Conjecture 0.2 is the result saying that each ω -categorical, characteristically simple p -group with NIP and having a non-algebraic, generically stable over \emptyset type is abelian (so, under all these assumptions, (iii) of Fact 0.1 cannot happen).

We also show that each infinite, ω -categorical, characteristically simple p -group with NIP has an infinite, definable abelian subgroup. Recall that the existence of an infinite, abelian subgroup is known for all infinite, ω -categorical groups (use the fact that such groups are locally finite together with [12, Corollary 2.5]). However, Plotkin [16] found examples of infinite, ω -categorical p -groups with no infinite, definable, abelian subgroup.

At the end of the paper, we observe that an ω -categorical group having at least one strongly regular type is abelian.

I am grateful to Dugald Macpherson for interesting discussions and suggestions.

1 Preliminaries

Recall that we say that a group G is solvable-by-finite [nilpotent-by-finite or abelian-by-finite] if it has a finite index (normal) subgroup which is solvable [nilpotent or abelian, respectively]. It is standard (see e.g. [11, Remark 2.5]) that if G is nilpotent-by-finite [abelian-by-finite], then it has a definable normal subgroup of finite index which is nilpotent [abelian, respectively]. If G is solvable-by-finite, it is not clear whether it has a definable, solvable subgroup of finite index (it has such a subgroup

if we additionally assume either icc on centralizers for all definable quotients of definable subgroups [10, Remark 3.3] or ω -categoricity).

Recall some basic notions from ring theory. In this paper, rings are associative, but they are not assumed to contain 1 or to be commutative. An element r of a ring R is nilpotent of nilexponent n if $r^n = 0$, and n is the smallest number with this property. The ring is nil [of nilexponent n] if every element is nilpotent [of nilexponent $\leq n$, and there is an element of nilexponent n]. The ring is nilpotent of class n if $r_1 \cdots r_n = 0$ for all $r_1, \dots, r_n \in R$, and n is the smallest number with this property. An element r is null if $rR = Rr = \{0\}$. The ring is null if all its elements are.

We say that a ring R is nilpotent-by-finite [null-by-finite] if it has a finite index ideal (equivalently subring by [13]) which is nilpotent [null, respectively]. By virtue of [11, Remark 2.7], this ideal can be chosen definable.

The Jacobson radical of a ring R is the collection of all elements of R satisfying the formula $\varphi(x) = \forall y \exists z (yx + z + zyx = 0)$. The ring R is semisimple if $J(R) = \{0\}$. It is always the case that $R/J(R)$ is semisimple.

Recall that a ring R is a subdirect product of rings R_i , $i \in I$, if there is a monomorphism of R into $\prod_{i \in I} R_i$ whose image projects onto each R_i . The following is [3, Corollary 1].

Fact 1.1 *If R is a semisimple, ω -categorical ring, then R is a subdirect product of complete matrix rings over finite fields. Moreover, only finitely many different matrix rings occur as subdirect factors.*

By [3, Lemma 1.3] and [4] we have

Fact 1.2 *If R is an ω -categorical ring, then $J(R)$ is nilpotent.*

So, in order to prove that an ω -categorical ring R satisfying some extra assumptions is nilpotent-by-finite, it is enough to show that the semisimple ring $R/J(R)$ is finite (here Fact 1.1 may be very helpful). We will use this approach in the proof of Theorem 2.1.

We will also use [11, Theorem 3.15].

Fact 1.3 *Suppose G is a solvable, ω -categorical group such that each ring interpretable in it is nilpotent-by-finite and each group H definable in it has a definable connected component H^0 (i.e. the smallest definable subgroup of finite index). Then G is nilpotent-by-finite.*

Now, we recall the relevant notions from model theory. Let T be a first order theory. We work in a monster model \mathfrak{C} of T .

Definition 1.4 *We say that T has the NIP if there is no formula $\varphi(x, y)$ and sequence $(a_i)_{i < \omega}$ such that for every $w \subseteq \omega$, there is b_w such that $\models \varphi(a_i, b_w)$ iff $i \in w$.*

The next fact is Theorem 1.0.5 of [18].

Fact 1.5 *If G is a group defined in a theory with NIP, then for each φ , there is some N such that the intersection of any finite family of φ -definable subgroups of G is an intersection of at most N members of the family.*

Let $p \in S(\mathfrak{C})$ be invariant over $A \subset \mathfrak{C}$. We say that $(a_i)_{i \in \omega}$ is a Morley sequence in p over A if $a_n \models p|Aa_{<n}$ for all n . It turns out that Morley sequences in p over A are indiscernible over A and they have the same type over A . If $\mathfrak{C}' \succ \mathfrak{C}$ is a bigger monster model, then the defining scheme of p determines a unique A -invariant extension $\tilde{p} \in S(\mathfrak{C}')$ of p . By a Morley sequence in p we mean a Morley sequence in \tilde{p} over \mathfrak{C} . Finally, $p^{(k)}$ (where $k \in \omega \cup \{\omega\}$) denotes the type over \mathfrak{C} of a Morley sequence in p of length k .

Recall from [15] that a global type $p \in S(\mathfrak{C})$ is said to be generically stable if, for some small A , it is A -invariant and for each Morley sequence $(a_i : i \in \omega)$ in p over A and each formula $\varphi(x)$ (with parameters from \mathfrak{C}), the set $\{i : \models \varphi(a_i)\}$ is either finite or co-finite. This definition does not depend on the choice of A over which p is invariant. We will say that p is generically stable over A to express that p is invariant over A and generically stable. Assuming NIP, there are various equivalent definitions of generic-stability (see [8, Proposition 3.2]). For us, one of them will be particularly important.

Fact 1.6 *Assume T has NIP, and $p \in S(\mathfrak{C})$ is A -invariant. Then, p is generically stable iff every/some Morley sequence in p over A is an indiscernible set over A .*

Now, we will briefly discuss fsg and generic stability. For more details on these and related notions see [7, 8].

Definition 1.7 *Let G be a group definable in \mathfrak{C} by a formula $G(x)$.*

(i) G has fsg (finitely satisfiable generics) if there is a global type p containing $G(x)$ and a model $M \prec \mathfrak{C}$, of cardinality less than the degree of saturation of \mathfrak{C} , such that for all g , gp is finitely satisfiable in M .

(ii) G is generically stable if G has fsg and some global generic type of G is generically stable.

We say that a group definable in a non-saturated model has one of the above properties if the group defined by the same formula in a monster model has it.

In general, generic stability of G is a strictly stronger notion than fsg, but it is easy to check that these notions agree when G^{00} (the smallest type-definable subgroup of bounded index) is definable and T has NIP (see [8, Section 6]). In the ω -categorical, NIP context, G^{00} is definable, and thus we get

Remark 1.8 *Assume G is a group definable in an ω -categorical structure with NIP. Then, it has fsg iff it is generically stable.*

At the end, we recall the notion of a strongly regular type from [15]. The geometric meaning of this notion is explained in [15].

Definition 1.9 *Let $p(x) \in S(\mathfrak{C})$ be non-algebraic. We say that $p(x)$ is strongly regular if, for some small A , it is A -invariant and for all $B \supseteq A$ and a from the sort of x , either $a \models p|B$ or $p|B \vdash p|Ba$.*

2 ω -categorical rings with NIP

In this section, we prove Conjecture 0.3.

Theorem 2.1 *Each ω -categorical ring with NIP is nilpotent-by-finite.*

Proof. By Fact 1.2, everything boils down to showing that a semisimple, ω -categorical ring R with NIP is finite. Suppose for a contradiction that R is infinite.

By ω -categoricity, the two sided ideals RrR , $r \in R$, are uniformly definable (because ω -categoricity implies that there is K such that every element of any RrR is the sum of at most K elements of the form r_1rr_2 for $r_1, r_2 \in R$). Thus, by NIP and Fact 1.5, there is $N \geq 1$ such that for all $n \in \omega$ and $r_0, \dots, r_n \in R$, there are $i_1, \dots, i_N \in \{0, \dots, n\}$ such that $Rr_0R \cap \dots \cap Rr_nR = Rr_{i_1}R \cap \dots \cap Rr_{i_N}R$.

Fact 1.1 tells us that R can be treated as a subring of the product $\prod_{i \in I} R_i$ of finite rings R_i with identity, which projects onto each R_i , and where there are only finitely many pairwise distinct rings among the R_i 's, $i \in I$. Let π_i be the projection onto the i th coordinate. For $i_0, \dots, i_n \in I$ and $r_j \in R_{i_j}$, we introduce the set

$$R_{i_0, \dots, i_n}^{r_0, \dots, r_n} = \left\{ r \in R : \bigwedge_{j=0}^n \pi_{i_j}(r) = r_j \right\}.$$

Using the assumption that R is infinite and R_i 's are finite, we see that for any $i_0, \dots, i_n \in I$,

$$R_{i_0, \dots, i_n}^{0, \dots, 0} \text{ is an infinite ideal of } R. \quad (*)$$

Claim 1 There are pairwise distinct $i_0, i_1, \dots \in I$, non-nilpotent elements $r_j \in R_{i_j}$, and elements $\eta_j \in R$ such that for all $n \in \omega$,

$$\eta_n \in R_{i_0, \dots, i_n}^{0, \dots, 0, r_n}.$$

Proof of Claim 1. Suppose $i_0, \dots, i_n, r_0, \dots, r_n$ and η_0, \dots, η_n have been chosen. Since R is semisimple, it has no non-trivial nil ideals. Thus, by (*), $R_{i_0, \dots, i_n}^{0, \dots, 0}$ contains a non-nilpotent element η_{n+1} . As there are only finitely many different R_i 's and they are all finite, we can find i_{n+1} different from i_0, \dots, i_n such that $r_{n+1} := \pi_{i_{n+1}}(\eta_{n+1}) \in R_{i_{n+1}}$ is non-nilpotent. \blacksquare

Claim 2 There are natural numbers $n(0) < \dots < n(N)$ such that the sets

$$R_{i_{n(0)}, \dots, i_{n(N)}}^{r_{n(0)}, 0, \dots, 0}, R_{i_{n(0)}, \dots, i_{n(N)}}^{0, r_{n(1)}, 0, \dots, 0}, \dots, R_{i_{n(0)}, \dots, i_{n(N)}}^{0, \dots, 0, r_{n(N)}}$$

are non-empty.

Before proving Claim 2, let us notice that it leads to a contradiction. Choose a_0, \dots, a_N from $R_{i_{n(0)}, \dots, i_{n(N)}}^{r_{n(0)}, 0, \dots, 0}, \dots, R_{i_{n(0)}, \dots, i_{n(N)}}^{0, \dots, 0, r_{n(N)}}$, respectively. Put $b_k = \sum_{l \neq k} a_l$ for $k = 0, \dots, N$. Then,

$$\pi_{i_{n(j)}}[Rb_0R \cap \dots \cap Rb_NR] = \{0\} \text{ for } j = 0, \dots, N. \quad (**)$$

On the other hand, $\prod_{k \neq j} b_k \in \bigcap_{k \neq j} Rb_k R$ for $j = 0, \dots, N$. We also have that $\pi_{i_{n(j)}}[\prod_{k \neq j} b_k] = r_{n(j)}^N \neq 0$ as $r_{n(j)}$ is non-nilpotent. So,

$$\pi_{i_{n(j)}}[\bigcap_{k \neq j} Rb_k R] \neq \{0\} \text{ for } j = 0, \dots, N. \quad (***)$$

By (**) and (***), $Rb_0 R \cap \dots \cap Rb_N R \neq \bigcap_{k \neq j} Rb_k R$ for all $j = 0, \dots, N$. This is a contradiction with the choice of N .

Proof of Claim 2. Let $c = \max_{i \in I} |R_i|$. Define recursively:

$$\begin{aligned} L_N &= c + 1, \\ L_{N-k} &= c^{L_N + \dots + L_{N-k+1}} + 1 \text{ for } k = 1, \dots, N-1. \end{aligned}$$

Put $L_0 = 0$.

Consider any $k \in \{0, \dots, N-1\}$. Suppose each natural number α from the closed interval $[L_{N-k-1} + \dots + L_0, L_{N-k} + \dots + L_0 - 1]$ has color

$$(\pi_{i_{L_{N-k} + \dots + L_1}}(\eta_\alpha), \dots, \pi_{i_{L_N + \dots + L_1}}(\eta_\alpha)) \in \prod_{j=L_{N-k} + \dots + L_1}^{L_N + \dots + L_1} R_{i_j}.$$

In this way, the natural numbers from the interval $[L_{N-k-1} + \dots + L_0, L_{N-k} + \dots + L_0 - 1]$ have been colored with at most $c^{L_N + \dots + L_{N-k+1}} = L_{N-k} - 1$ colors. Since there are L_{N-k} such numbers, we can find natural numbers $n(N-k-1) < n'(N-k-1)$ from $[L_{N-k-1} + \dots + L_0, L_{N-k} + \dots + L_0 - 1]$ with the same color. Put $a_{N-k-1} = \eta_{n(N-k-1)} - \eta_{n'(N-k-1)}$. Then, $\pi_{i_j}(a_{N-k-1}) = 0$ for all $j \in [L_N + \dots + L_{N-k}, L_N + \dots + L_1]$. Moreover, from the choice of η_m 's, we see that $\pi_{i_j}(a_{N-k-1}) = 0$ for all $j < L_{N-k-1} + \dots + L_0$. Putting additionally $n(N) = L_N + \dots + L_1$, we get a sequence $n(0) < \dots < n(N)$ which satisfies the conclusion of the claim. \blacksquare

3 ω -categorical groups with NIP

In this section, we investigate the structure of ω -categorical groups with NIP. First, we make some observations on characteristically simple groups in this context. Then, we prove Conjecture 0.2 under the additional assumptions of fsg or of the existence of a strongly regular type.

As was mentioned in the introduction, each ω -categorical group is locally finite, and so, if it is infinite, it has an infinite, abelian subgroup [12, Corollary 2.5]. However, [16] yields examples of infinite, ω -categorical p -groups with no infinite, definable, abelian subgroup.

Proposition 3.1 *Let p be a prime number. Then every infinite, characteristically simple, ω -categorical p -group G with NIP has an infinite, definable, abelian subgroup.*

Proof. For any $x_0, \dots, x_n \in G$, $\langle x_0, \dots, x_n \rangle$ is a finite p -group, and so $C(x_0) \cap \dots \cap C(x_n) \neq \{e\}$. Hence, if there were $x_0, \dots, x_n \in G$ with $C(x_0) \cap \dots \cap C(x_n)$ finite, we would find $x_{n+1}, \dots, x_m \in G$ such that $Z(G) = C(x_0) \cap \dots \cap C(x_m) \neq \{e\}$, which would contradict the characteristic simplicity of G . So, we have proved the following

Claim For any $n \in \omega$ and $x_0, \dots, x_n \in G$, the intersection $C(x_0) \cap \dots \cap C(x_n)$ is infinite.

By the Claim, we can choose a sequence $(x_n)_{n \in \omega}$ of pairwise distinct elements of G such that $x_{n+1} \in C(x_0) \cap \dots \cap C(x_n)$ for all $n \in \omega$. Put $G_n = C(x_0) \cap \dots \cap C(x_n)$. Then, $x_0, \dots, x_n \in Z(G_n)$. Hence, $|Z(G_n)| > n$. On the other hand, by NIP and Fact 1.5, there is N such that for any $y_0, \dots, y_n \in G$, there are $i_1, \dots, i_N \in \{0, \dots, n\}$ with $C(y_0) \cap \dots \cap C(y_n) = C(y_{i_1}) \cap \dots \cap C(y_{i_N})$. Thus, $G_n = C(x_{i_1^n}) \cap \dots \cap C(x_{i_N^n})$ for some $i_1^n, \dots, i_N^n \in \{0, \dots, n\}$. Since by ω -categoricity the set $\{tp(x_{i_1^n}, \dots, x_{i_N^n}) : n \in \omega\}$ is finite, there is $n \in \omega$ such that $Z(G_n)$ is infinite. ■

The next proposition uses notation from Fact 0.1.

Proposition 3.2 *For any non-abelian, finite, simple group F , neither $B(F)$ nor $B^-(F)$ have NIP.*

Proof. Let C_0, C_1, \dots be disjoint clopen subsets of the Cantor set \mathcal{C} not containing x_0 . Choose $g \in F \setminus Z(F)$. Define a sequence $(f_i)_{i \in \omega}$ of elements of $B^-(F)$ by:

$$f_i(\eta) = \begin{cases} g & \text{if } \eta \in C_i, \\ e & \text{if } \eta \notin C_i. \end{cases}$$

Now, suppose for a contradiction that $B(F)$ has NIP (the case when $B^-(F)$ has NIP is almost the same). Using NIP and Fact 1.5, and reordering C_i 's if necessary, we can find N such that $C_{B(F)}(f_0) \cap \dots \cap C_{B(F)}(f_N) = C_{B(F)}(f_0) \cap \dots \cap C_{B(F)}(f_{N-1})$.

Take $h \in F \setminus C(g)$ and define $f \in B^-(F)$ by:

$$f(\eta) = \begin{cases} h & \text{if } \eta \in C_N, \\ e & \text{if } \eta \notin C_N. \end{cases}$$

Then, we see that $f \in C_{B(F)}(f_0) \cap \dots \cap C_{B(F)}(f_{N-1}) \setminus C_{B(F)}(f_0) \cap \dots \cap C_{B(F)}(f_N)$, a contradiction. ■

Proposition 3.3 *Let p be a prime number. Let G be a characteristically simple p -group interpretable over \emptyset in an ω -categorical structure with NIP. Assume that G has a global, non-algebraic type q which is generically stable over \emptyset . Then G is abelian.*

Proof. Assume for simplicity that G is the ambient structure. Wlog G is a monster model. Let $(a_i)_{i \in \omega}$ be a Morley sequence in q over \emptyset . By NIP and Fact 1.5, there is N such that for any m , $C(a_0) \cap \dots \cap C(a_m) = C(a_{i_1}) \cap \dots \cap C(a_{i_N})$ for some $i_1, \dots, i_N \in \{0, \dots, m\}$. But, using Fact 1.6, $(a_i)_{i \in \omega}$ is an indiscernible set over

\emptyset . This implies that for any $m \geq N - 1$ and $0 \leq i_1 < \dots < i_N \leq m$, one has $C(a_0) \cap \dots \cap C(a_m) = C(a_{i_1}) \cap \dots \cap C(a_{i_N})$.

Consider any $(b_0, b_1, \dots) \models q^{(\omega)} | \emptyset$. Then, there is a sequence $(c_i)_{i \in \omega}$ such that the sequences $(a_i : i \in \omega) \frown (c_i : i \in \omega)$ and $(b_i : i \in \omega) \frown (c_i : i \in \omega)$ are indiscernible over \emptyset . Thus, by the last paragraph, $\bigcap_{i \in \omega} C(a_i) = C(c_0) \cap \dots \cap C(c_{N-1}) = \bigcap_{i \in \omega} C(b_i)$. But, $\langle c_0, \dots, c_{N-1} \rangle$ is a finite p -group, which implies that $C(c_0) \cap \dots \cap C(c_{N-1}) \neq \{e\}$. We conclude that $\bigcap_{i \in \omega} C(a_i)$ is a non-trivial, \emptyset -invariant (so \emptyset -definable) subgroup. Since G is characteristically simple, we get $\bigcap_{i \in \omega} C(a_i) = G$, which implies that $Z(G) \neq \{e\}$, and so $G = Z(G)$ once again by the characteristic simplicity of G . ■

Theorem 3.4 *Each ω -categorical group G with NIP and fsg is nilpotent-by-finite.*

Proof. We can assume that G is infinite and it is a monster model. In fact, we will work in a slightly more general context, where G is a group with fsg which is interpretable (over \emptyset) in an ω -categorical monster model $\mathfrak{C} = \mathfrak{C}^{eq}$ satisfying NIP. Since G^{00} is \emptyset -invariant, it is \emptyset -definable by ω -categoricity. Thus, we can assume that $G = G^{00}$. Then, G has a unique global generic type [5, Proposition 0.26], which must be \emptyset -invariant.

By ω -categoricity, there is a normal series $\{e\} = G_0 \leq G_1 \leq \dots \leq G_n = G$ of \emptyset -definable (in \mathfrak{C}) subgroups of G such that each G_{i+1}/G_i is characteristically simple in the sense of (G, \mathfrak{C}) , i.e. it does not have non-trivial, proper, subgroups normal in G/G_i and invariant under $Aut(\mathfrak{C})$. In particular, each G_{i+1}/G_i is a characteristically simple group. Indeed, it is clear that any characteristic subgroup of G_{i+1}/G_i is invariant under $Aut(\mathfrak{C})$; the fact that such a subgroup is normal in G/G_i follows from the fact that inner automorphisms of G induce automorphisms of G_{i+1}/G_i .

Our proof will be by induction on n (where n is a number for which there exists a sequence as in the last paragraph). Consider the case $n = 1$. Then, $G = G_1$ is an infinite, characteristically simple group. Proposition 3.2 eliminates the possibility that a countable elementary substructure of G is as in point (ii) of Fact 0.1. Proposition 3.3 together with Remark 1.8 and the \emptyset -invariance of the unique generic type of G eliminate the possibility from point (iii) of Fact 0.1. Thus, G must be abelian.

We turn to the induction step, where we assume that $n \geq 2$. First, notice that ω -categoricity implies that the upper central series $Z_0(G) \leq Z_1(G) \leq Z_2(G) \leq \dots$ stabilizes after finitely many, say m , steps. So, replacing G by $G/Z_m(G)$, we can assume that G is centerless. Such a replacement is possible, because the normal series $G_0 Z_m(G)/Z_m(G) \leq G_1 Z_m(G)/Z_m(G) \leq \dots \leq G_n Z_m(G)/Z_m(G) = G/Z_m(G)$ consists of groups which are \emptyset -definable in \mathfrak{C} , and it is easy to check that each quotient $(G_{i+1} Z_m(G)/Z_m(G))/(G_i Z_m(G)/Z_m(G))$ is characteristically simple in the sense of $(G/Z_m(G), \mathfrak{C})$.

Since $n \geq 2$, by the induction hypothesis, we can assume that G_1 is a non-trivial, proper subgroup of G . Since G/G_1 also has fsg, using the induction hypothesis, we get that G/G_1 is nilpotent-by-finite (notice that since we consider \emptyset -definability in \mathfrak{C} , the normal series $G_1/G_1 \leq G_2/G_1 \leq \dots \leq G_n/G_1 = G/G_1$ allows us to use the induction hypothesis). So, it is nilpotent, because the Fitting subgroup of G/G_1 is a

nilpotent subgroup of finite index which is \emptyset -definable and so equal to G/G_1 by the connectedness of G/G_1 .

Let q be the unique global generic type of G , and $(g_i)_{i \in \omega}$ be a Morley sequence in q over \emptyset . Since G/G_1 is nilpotent, there is a minimal k such that the iterated commutator $[g_{k-1}, [g_{k-2}, \dots, [g_1, g_0] \dots]] \in G_1$. Since g_0 is generic over \emptyset and G/G_1 is infinite, we see that $k \geq 2$. Define

$$h_i = [g_{ik+k-1}, [g_{ik+k-2}, \dots, [g_{ik+1}, g_{ik}] \dots]]$$

for $i \in \omega$. Let $(g'_i)_{i \in \omega}$ be a Morley sequence in q over \mathfrak{C} . Put

$$h'_i = [g'_{ik+k-1}, [g'_{ik+k-2}, \dots, [g'_{ik+1}, g'_{ik}] \dots]]$$

for $i \in \omega$. Since $tp(g'_{k-1}, \dots, g'_0/\mathfrak{C}) = q^{(k)}$ is invariant over \emptyset , the type $r := tp(h'_0/\mathfrak{C})$ is also invariant over \emptyset . Moreover, $(h_i)_{i \in \omega}$ is a Morley sequence in r over \emptyset . By the generic stability of q and Fact 1.6, the sequence $(g_i)_{i \in \omega}$ is an indiscernible set, and so $(h_i)_{i \in \omega}$ is an indiscernible set as well. We conclude that r is generically stable over \emptyset .

We claim that r is non-algebraic. To see this, let us define

$$a_j := [g'_{k-1+j}, [g'_{k-2}, \dots, [g'_1, g'_0] \dots]]$$

for $j \in \omega$. We see that $a_j \models r$. Moreover, $a_{j_1} \neq a_{j_2}$ whenever $j_1 < j_2$. Indeed, if this is not the case, then for some $j_1 < j_2$, we have that $g'_{j_2} g'^{-1}_{j_1} \in C(h)$, where $h = [g'_{k-2}, \dots, [g'_1, g'_0] \dots]$. By the minimality of k , $h \neq e$. But, g'_{j_2} is generic over g'_{j_1}, h , and so $g'_{j_2} g'^{-1}_{j_1}$ is generic over h . Thus, $[G(\mathfrak{C}') : C(h)] < \omega$, where \mathfrak{C}' is a monster model containing \mathfrak{C} and $(g'_i)_{i \in \omega}$. This implies $h \in Z(G(\mathfrak{C}')) = \{e\}$, a contradiction.

We have proved that r is a global, non-algebraic type of G_1 which is generically stable over \emptyset (in particular, G_1 is infinite). So, by Proposition 3.3 together with Proposition 3.2 and Fact 0.1, G_1 is abelian. Hence, G is solvable.

By Theorem 2.1 and Fact 1.3, we conclude that G is nilpotent-by-finite. ■

Dugald Macpherson told me an alternative ending of the above proof, i.e. an alternative proof of the fact that a solvable, ω -categorical group with NIP is nilpotent-by-finite. Namely, by [2], we know that each countable, solvable, ω -categorical group which is not nilpotent-by-finite interprets the countable, atomless Boolean algebra. So, it remains to show that this algebra does not have NIP, which is an easy exercise.

Now, we will drop the NIP and fsg assumption, and instead we will assume the existence of a strongly regular type. Recall the following question from [15].

Question 3.5 *Suppose G is a group with at least one strongly regular type. Does it imply that G is abelian?*

Proposition 3.6 *If G is any group with at least one strongly regular type, then all non-central elements of G are conjugated.*

Proof. Taking an elementary extension of G , we can assume that there is a global type p whose strong regularity is witnessed over G . Consider any non-central element $a \in G$. Take $b \models p|G$.

Notice that if $a^b \models p|G$, then the formula defining the conjugacy class of a belongs to $p|G$. Thus, all elements $a \in G$ for which $a^b \models p|G$ are in one conjugacy class. So, it remains to show that the assumption $a^b \not\models p|G$ leads to contradiction.

This assumption and the strong regularity of p over G imply that $tp(b/G) \vdash tp(b/a^b, G)$. Thus, there is a formula $\varphi(x, y)$ (without parameters) and $g \in G^n$ such that $b \models \varphi(x, g)$ and $\models (\varphi(x, g) \rightarrow a^x = a^b)$. So, there is $c \in G$ such that $a^c = a^b$, and hence $b \in C(a)c$. This means that $p|G \vdash 'x \in C(a)c'$.

Consider two distinct realizations g_1 and g_2 of $p|G$ (they exist because p is non-algebraic).

Case 1 $c \in C(a)$.

Then, $p|G \vdash 'x \in C(a)'$, so $g_1 \in C(a)$. Take $h \notin C(a)$. Then, $hg_1 \notin C(a)$. Thus, by the strong regularity of p over G , we conclude that $tp(g_1/G, h, hg_1) = p|G, h, hg_1 = tp(g_2/G, h, hg_1)$. But, the formula $x = h^{-1}hg_1$ belongs to $tp(g_1/G, h, hg_1)$ and does not belong to $tp(g_2/G, h, hg_1)$, a contradiction.

Case 2 $c \notin C(a)$.

Since $g_1 \in C(a)c$, we have $g_1c^{-1} \in C(a)$, and so $g_1c^{-1} \notin C(a)c$. Hence, by the strong regularity of p , we get $tp(g_1/G, g_1c^{-1}) = p|G, g_1c^{-1} = tp(g_2/G, g_1c^{-1})$. But, the formula $x = g_1c^{-1}c$ belongs to $tp(g_1/G, g_1c^{-1})$ and does not belong to $tp(g_2/G, g_1c^{-1})$, a contradiction. ■

Corollary 3.7 *If G is a group with at least one strongly regular type, then all non-central elements of G have infinite order. In particular, an ω -categorical group with at least one strongly regular type is abelian.*

Proof. This is a standard argument. We can assume that $G \neq Z(G)$. Suppose for a contradiction that there is a non-central element of finite order. By the last proposition, $G/Z(G)$ has one non-trivial conjugacy class. So, all non-trivial elements of $G/Z(G)$ have the same order, which must be a prime number p . If $p = 2$, then $G/Z(G)$ is abelian, so $[G : Z(G)] \leq 2$, which implies $G = Z(G)$, a contradiction. Now, we assume that p is odd. Take a nontrivial $g \in G/Z(G)$. Then, there is $h \in G/Z(G)$ such that $h^{-1}gh = g^{-1}$. So, $g \in C(h^2) \setminus C(h)$. Finally we get

$$C(h) = C(h^{2^{p-1}}) \supsetneq C(h^{2^{p-2}}) \supsetneq \cdots \supsetneq C(h^2) \supsetneq C(h),$$

which is impossible. ■

References

- [1] A. Apps, *On the structure of \aleph_0 -categorical groups*, Journal of Algebra 81, 320-339, 1983.

- [2] R. Archer, D. Macpherson, *Soluble, ω -categorical groups*, Math. Proc. Camb. Phil. Soc.121, 219-227, 1997.
- [3] J. Baldwin, B. Rose, *\aleph_0 -categoricity and stability of rings*, Journal of Algebra 45, 1-16, 1977.
- [4] G. Cherlin, *On \aleph_0 -categorical nilrings. II*, Journal of Symbolic Logic 45, 291-301, 1980.
- [5] C. Ealy, K. Krupiński, A. Pillay, *Superrosy dependent groups having finitely satisfiable generics*, Annals of Pure and Applied Logic 151, 1-21, 2008.
- [6] D. Evans, F. Wagner, *Supersimple ω -categorical groups and theories*, Journal of Symbolic Logic 65, 767-776, 2000.
- [7] E. Hrushovski, Y. Peterzil, A. Pillay, *Groups, measures and the NIP*, Journal of the AMS 21, 563-596, 2008.
- [8] E. Hrushovski, A. Pillay, *On NIP and invariant measures*, preprint, 2009.
- [9] O. Kegel, B. Wehrfritz, *Locally Finite Groups*, North-Holland, The Netherlands, 1973.
- [10] K. Krupiński, *Fields interpretable in rosy theories*, Israel Journal of Mathematics 175, 421-444, 2010.
- [11] K. Krupiński, *On relationships between algebraic properties of groups and rings in some model-theoretic contexts*, submitted.
- [12] K. Krupiński, F. Wagner, *Small profinite groups and rings*, Journal of Algebra 306, 494-506, 2006.
- [13] J. Lewin, *Subrings of finite index in finitely generated rings*, Journal of Algebra 5, 84-88, 1967.
- [14] H. D. Macpherson, *Absolutely ubiquitous structures and \aleph_0 -categorical groups*, Quart. J. Math. Oxford (2) 39, 483-500, 1988.
- [15] A. Pillay, P. Tanović, *Generic stability, regularity, and quasi-minimality*, preprint, 2009.
- [16] J. M. Plotkin, *ZF and locally finite groups*, Zeitschrift f. Math. Logik and Grundlagen der Mathematik 27, 375-379, 1981.
- [17] B. Poizat, *Stable groups*, American Mathematical Society, Providence, 2001.
- [18] F. Wagner, *Stable groups*, London Math. Soc. Lecture Notes Series 240, Cambridge University Press, UK, 1997.

- [19] J. Wilson, *The algebraic structure of \aleph_0 -categorical groups*, in Groups-St. Andrews, Ed. C. M. Campbell, E. F. Robertson, London Math. Soc. Lecture Notes 71, Cambridge, 345-358, 1981.

Krzysztof Krupiński
Instytut Matematyczny Uniwersytetu Wrocławskiego
pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland.
e-mail: kkrup@math.uni.wroc.pl