

G -linear sets and torsion points in definably compact groups

Margarita Otero*
Universidad Autónoma de Madrid

Ya'acov Peterzil
University of Haifa

August 3, 2007

Abstract

Let G be a definably compact group in an o-minimal expansion of a real closed field. We prove that if $\dim(G \setminus X) < \dim G$ for some definable $X \subseteq G$ then X contains a torsion point of G . Along the way we develop a general theory for so-called G -linear sets, and investigate definable sets which contain abstract subgroups of G .

Keywords: o-minimality, definable group, torsion point.

Mathematics Subject Classification 2000: 03C64

1 Introduction

We prove:

Theorem 1.1. *Let G be a definably compact group in an o-minimal expansion of a real closed field. If $X \subseteq G$ is a definable large set (i.e. $\dim(G \setminus X) < \dim G$) then X contains a torsion point of G .*

This is a weak approximation to the conjecture that every generic set in a definably compact group contains a torsion point, a conjecture which itself follows from the “compact domination conjecture” from [4].

The proof starts with the abelian case, where we first develop, in o-minimal expansions of an ordered group, a notion of a coset of a definable local subgroup (a G -linear set). We then use tools from Algebraic Topology, in expansions of real closed fields, to conclude the theorem.

*Partially supported by GEOR MTM2005-02568 and Grupos UCM 910444

2 G -linear sets

In this section \mathcal{M} will denote an o-minimal expansion of an ordered group. By definable we mean definable in \mathcal{M} . G will denote a definable abelian group. All topological concepts are with respect to the group topology such G is equipped with. (We believe that most, if not all, of the work below can be developed in an arbitrary definable group, but for simplicity we limit ourselves to the commutative setting.)

Definition 2.1. 1) Given two sets $X, Y \subseteq G$ and $a \in G$, we say that X and Y have the same germ at a , in notation $X =_a Y$, if there exists an open neighborhood U of a such that $X \cap U = Y \cap U$. We say that the germ of X at a is contained in the germ of Y at a , in notation $X \subseteq_a Y$, if there exists an open $U \ni a$ such that $X \cap U \subseteq Y \cap U$.

2) Given $X \subseteq G$ and $g, h \in G$, we say that the germ of X at g is G -equivalent to the germ of X at h if $X - g =_0 X - h$. (Note that if X is a definable set then we obtain in this way a definable equivalence relation on G .)

Definition 2.2. Let X be a definable subset of G .

1) X is called G -linear if for every $g, h \in X$ we have $X - g =_0 X - h$. Given $g \in X$, we say that X is locally G -linear at g if there exists an open $U \ni g$ such that for every $h \in U \cap X$ we have $X - g =_0 X - h$.

Two G -linear sets $X, Y \subseteq G$ are called G -equivalent if for every $g \in X$, $h \in Y$, we have $X - g =_0 Y - h$.

2) Let X be a G -linear set. A definable $Y \subseteq G$ is called a G -subset of X if for every $h \in Y$ and $g \in X$, we have $Y - h \subseteq_0 X - g$.

Example 2.3. Let G be a definable abelian group and H a definable subgroup. If X is a relatively open subset of a coset of H then X is G -linear. If Y is any definable subset of a coset of H then Y is a G -linear subset of X .

We observe without a proof (we will not be using it): If \mathcal{M} expands a real closed field then Y is a G -subset of the G -linear set X if and only if at every smooth point $y \in Y$, the tangent space of Y at y is a subset of the tangent space of X at any of its points.

Lemma 2.4. If X is definably connected and locally G -linear at every $g \in X$ then X is G -linear.

Proof. Because X is locally G -linear at every point the G -equivalence class of every $g \in X$ is relatively open in X . But then it is also relatively closed

(since its complement is a union of open sets), so by definable connectedness there exists only one G -equivalence class, so X is G -linear. \square

Definition 2.5. *Given $X \subseteq G$, we let*

$$X_{lin} = \{g \in X : X \text{ is locally } G\text{-linear at } g\}.$$

Notice that X_{lin} is a relatively open (possibly empty) subset of X . By the last lemma, every definably connected component of X_{lin} is a G -linear set, but X_{lin} itself might not be G -linear (E.g., if $X = H_1 \cup H_2$ is the union of two definable subgroups of G , none containing the other, then $X_{lin} = X \setminus (H_1 \cap H_2)$ is not a G -linear set). It is easy to see that X_{lin} is itself G -linear if and only if all its definably connected components are G -equivalent.

Notation We denote by X_{lin}^{max} the union of all components of X_{lin} of maximal dimension.

Infinitesimals

It might be easier here to use the language of infinitesimals: We move to an $|M|^+$ -saturated elementary extension \mathcal{N} of \mathcal{M} . For $g \in G(\mathcal{M})$, we denote by ν_g the intersection of all \mathcal{M} -definable open neighborhoods of g (with respect to the group topology) in the structure \mathcal{N} . For $X \subseteq G$, we write $\nu_g(X) = \nu_g \cap X$. Notice that for $X, Y \subseteq G$ and $g, h \in G$, $\nu_g(X) = \nu_g(Y)$ iff $X =_g Y$, and $\nu_g(X) - g = \nu_h(X) - h$ iff $X - g =_0 X - h$.

We will be using the following simple observation: If X, Y are A -definable sets, and x is generic in X and in Y over A then $\nu_x(X) \subseteq Y$.

Lemma 2.6. *X is locally G -linear at $g \in X$ if and only if $\nu_g(X) - g$ is a subgroup of G . In particular, X is a C^0 -manifold near g .*

Proof. : If $\nu_g(X) - g$ is a subgroup then for every $h \in \nu_g(X)$, we have $h - g + \nu_g(X) = \nu_g(X)$. It easily follows that X is locally G -linear at g .

Conversely, assume that X is locally G -linear at g . We need to prove: For all $h_1, h_2 \in X$ sufficiently close to g we have $(h_1 - g) - (h_2 - g) = h' - g$, for some $h' \in X$, or equivalently, $g + (h_1 - h_2) \in X$.

We pick $h \in X$ close to g and generic over g such that $X - g =_0 X - h$. We take $U \ni 0$, definable over parameters independent of g, h , such that $(X - g) \cap U = (X - h) \cap U$. Because of the genericity of h , there exists a neighborhood $V \ni h$, which we may assume is contained in $h + U$, such that for all $h' \in X \cap V$,

$$(X - h) \cap U = (X - g) \cap U = (X - h') \cap U.$$

By adding h to both sides of the last equation, we obtain: For all $h' \in X \cap V$, $X \cap (h + U) = (X - h' + h) \cap (h + U)$, hence (since $V \subseteq (h + U)$) also for all $h' \in X \cap V$,

$$X \cap V = (X - h' + h) \cap V.$$

In particular, if $h'', h' \in X$ are sufficiently close to h (so that $h'' - h' + h \in V$) then we have $(h'' - h') + h$ in X . It follows, as was pointed out above, that $\nu_h(X) - h$ is a subgroup of G . Because $\nu_g(X) - g = \nu_h(X) - h$, it also follows that $\nu_g(X) - g$ is a subgroup of G .

Because the germ of X at g is G -equivalent to the germ of X at some generic point near g , it follows that X is a C^0 -manifold near g . \square

Notation. For a G -linear X , we denote by $\nu(X)$ the group $\nu_x(X) - x$, for some (all) $x \in X$. Notice that Y is a G -subset of X iff $\nu_h(Y) - h$ is a subset of the group $\nu(X)$, for every $h \in Y$. Note also that two G -linear sets X_1, X_2 are G -equivalent if and only if $\nu(X_1) = \nu(X_2)$.

Lemma 2.7. *Let $X, Y \subseteq G$ be definable sets of dimension d . If $\dim(X + Y) = d$ then X_{lin} is large in X , Y_{lin} large in Y and $X_{lin}^{max}, Y_{lin}^{max}$ are G -linear sets that are G -equivalent.*

Proof. We assume that X and Y are \emptyset -definable and write $Z = X + Y$. Take g generic in X over \emptyset and h generic in Y over g . Then $g + h$ is generic in Z over each g and h and hence for every two $x, y \in \{g, h, g + h\}$ we have $\dim(x/y) = d$.

For every \emptyset -definable neighborhood U of g , we have $(U \cap X) + h \subseteq Z$. Moreover $g + h$ is generic in $(U \cap X) + h$ over h and therefore $(U \cap X) + h$ contains $\nu_{g+h}(Z)$. It follows that $\nu_g(X) + h \supseteq \nu_{g+h}(Z)$. By symmetry we may conclude that $\nu_g(X) + h = \nu_{g+h}(Z)$, or equivalently, $\nu_g(X) - g = \nu_{g+h}(Z) - (g + h)$.

In the very same way we see that $\nu_h(Y) - h = \nu_{g+h}(Z) - (g + h)$. Hence, we showed that for *every* generic $g \in X$, $h \in Y$ which are independent from each other we have $\nu_g(X) - g = \nu_h(Y) - h$.

Fixing a generic $h \in Y$, this implies that $\nu_{g'}(X)$ is constant as g' varies over all elements g' which are generic in X over h . In particular, X_{lin} contains all those elements g' and moreover all the components of X_{lin} of dimension d are G -equivalent. The same argument shows that all components of Y_{lin} of dimension d are G -equivalent to each other and that X_{lin}^{max} is G -equivalent to Y_{lin}^{max} . \square

Corollary 2.8. *Let $X, Y, Z \subseteq G$ be definable sets of dimension d . Assume that for every g in a large set $X_0 \subseteq X$, we have $\dim((g+Y) \cap Z) = d$. Then X_{lin} is large in X .*

Proof. Without loss of generality, X, Y, Z, X_0 are \emptyset -definable, and hence every generic element $g \in X$ belongs to X_0 . For every such $g \in X_0$ there exists $c \in (g+Y) \cap Z$ which is generic in Z over g . It follows that $h = c - g$ is in Y and, as before, the dimension of any two of $\{g, h, c\}$ is $2d$. We can now find open neighborhoods V of h and W of c , defined over independent parameters, such that $g + (V \cap Y) \subseteq (W \cap Z)$. Because of the genericity of g there is a neighborhood U of g such that $(U \cap X) + (V \cap Y) \subseteq (W \cap Z)$. In particular, $\dim((U \cap X) + (V \cap Y)) = d$. We can therefore apply Lemma 2.7 to $U \cap X$ and $V \cap Y$ and conclude that X is locally G -linear at g . \square

Note that we cannot conclude, under the assumption of the last Lemma, that all components of X_{lin}^{max} are G -equivalent.

Definition 2.9. *If U is an open symmetric ($U = -U$) neighborhood of 0 and $Y \subseteq G$ a definable set, we say that $h_1, h_2 \in Y$ are U -connected in Y if $h_2 \in h_1 + U$ and there exists a definable path in Y connecting h_1 and h_2 , which is contained in $h_1 + U$.*

Lemma 2.10. *Let X be G -linear and Y a G -subset of X . Take $g \in X$, and assume that U is an open symmetric neighbourhood of 0 such that $(g+U) \cap X$ is relatively closed in $g+U$. Then, for any $h_1, h_2 \in Y$ that are U -connected in Y , we have $g + (h_1 - h_2) \in X$*

Proof. Because h_1, h_2 are U -connected in Y , there exists a definable path $\gamma : [a, b] \rightarrow Y$ such that $\gamma(a) = h_1, \gamma(b) = h_2$ and for every $t \in [a, b]$, $\gamma(t) - h_1 \in U$. Consider the set T of all $t \in [a, b]$ such that $g + (h_1 - \gamma(t)) \in X$, for all $t' \leq t$.

We claim that T is both open and closed (with respect to the order topology in M) in $[a, b]$. Indeed, because $(g+U) \cap X$ is relatively closed in $g+U$, the set T is closed in $[a, b]$.

To see that it is open, assume that $t_0 \in T$. In particular, $k = g + (h_1 - \gamma(t_0))$ is in X . For $t' \in [a, b]$ close to t_0 we have $(\gamma(t_0) - \gamma(t')) \in X - k$ (because $\nu_{\gamma(t_0)}(Y) - \gamma(t_0) \subseteq \nu_k(X) - k = \nu(X)$ is a subgroup of G), therefore $t' \in T$. It follows that T is open and closed in $[a, b]$, hence $T = [a, b]$ and therefore $g + (h_1 - h_2) \in X$. \square

The following technical lemma ensures that we can extend every G -linear set and every G -subset of a G -linear set beyond its frontier (where the frontier of X is $Cl(X) \setminus X$).

Lemma 2.11. *Assume that X is G -linear and Y is G -subset of X . Then $Cl(Y)$ is also a G -subset of X .*

Proof. Fix h in $Cl(Y) \setminus Y$. We first prove that $\nu_h(Y) - h \subseteq \nu(X)$. Namely we show: For all $h' \in Y$ sufficiently close to h and every $g \in X$ we have $g + h' - h \in X$.

If the above fails then we have a curve $\gamma : [a, b] \rightarrow Y$, with $\lim_{t \rightarrow b} \gamma(t) = h$ such that for all $t \in [a, b)$, $g + \gamma(t) - h \notin X$. Fix U an open symmetric neighborhood of 0 such that $X \cap (g + U)$ is relatively closed in $g + U$. By choosing $\gamma(a)$ sufficiently close to h , we may assume that for every $t \in [a, b]$, we have $\gamma(t) \in \gamma(a) + U$ (so in particular, $h \in \gamma(a) + U$). It follows that for every $t \in (a, b)$, we have $\gamma(a)$ and $\gamma(t)$ are U -connected (as witnessed by γ) and therefore, by Lemma 2.10, $g + \gamma(a) - \gamma(t) \in (g + U) \cap X$. Because $X \cap (g + U)$ is closed in $g + U$, we may take t to be b and conclude that $g + \gamma(a) - \gamma(b) = g + \gamma(a) - h \in X$, contradicting our assumption.

We therefore showed that for all $h \in Cl(Y) \setminus Y$, $\nu_h(Y) - h \subseteq \nu(X)$. By our assumption on Y this is true for every $h \in Cl(Y)$. Because X is locally closed we may take the closure on the left and conclude that $\nu_h(Cl(Y)) - h \subseteq \nu(X)$ for every $h \in Cl(Y)$, hence $Cl(Y)$ is a G -subset of X . \square

Lemma 2.12. *Let $X \subseteq G$ be a definable set. Assume that*

- (i) X_{lin} is large in X (i.e. $\dim(X \setminus X_{lin}) < \dim(X)$) and
- (ii) for every $h \in X$ and $g \in X_{lin}$, $X - g \subseteq_0 X - h$.

Then X is G -linear.

Proof. Assume that $\dim(X) = d$. First notice that by (ii), all the components of X_{lin} are G -equivalent to each other and therefore X_{lin} is itself a G -linear set of dimension d .

Because of (ii) the local dimension of X at every point is d and therefore, by (i), X_{lin} is dense in X . By Lemma 2.11, X is a G -subset of X_{lin} .

Given any $h \in X$, we have

$$\nu_h(X) - h \subseteq \nu(X_{lin}) \subseteq \nu_h(X) - h.$$

(the left inclusion follows from the fact that X is a G -subset of X_{lin} while the right one is just assumption (ii)).

It follows that X is locally G -linear at h hence X is G -linear. \square

Before the next lemma we make a small observation.

Remark 2.13. *Let $Z, W \subseteq G$ be definable sets and let $f : Z \rightarrow W$ be a definable continuous map (all are 0-definable). Then for every $w \in f(Z)$ that*

is locally generic in W and $z \in f^{-1}(w)$, we have $\nu_w(W) = f(\nu_z(Z))$. (By “locally generic” we mean that for some open $V \ni w$, we have $\dim(w/\emptyset) = \dim(V \cap W)$).

Proof. Fix V as above and let $U \subseteq G$ be a definable open neighborhood of z such that w is still generic in $V \cap W$ over the parameters (say A) defining U . Then $\dim(w/A) = \dim(w) = \dim f(U \cap Z)$, $w \in f(U \cap Z)$ and therefore $f(U \cap Z)$ and $V \cap W$ have the same germ at w . In particular, $\nu_w(W) = \nu_w(V \cap W) \subseteq f(U \cap Z)$. Because this is true for every U neighborhood of z , it follows that $\nu_w(W) \subseteq f(\nu_z(Z))$. The converse follows by continuity. \square

Lemma 2.14. *Let X and Y be two G -linear sets. Then:*

(i) $X + Y$ is G -linear and we have

$$\nu(X + Y) = \nu(X) + \nu(Y).$$

In particular, if X and Y are G -equivalent G -linear sets then $X + Y$ is G -equivalent to them as well.

(ii) *If X and Y are G -equivalent then $X \cup Y$ is a G -linear set and G -equivalent to X and Y .*

Proof. (i) First, notice that, by continuity, for any $x \in X$, $y \in Y$, and $z = x + y$ we have $\nu_x(X) + \nu_y(Y) \subseteq \nu_z(X + Y)$. It follows that for all $z \in X + Y$, we have $\nu(X) + \nu(Y) \subseteq \nu_z(X + Y) - z$.

By Remark 2.13, if $z = x + y$ is a generic element of $X + Y$ then $\nu_z(X + Y) - z = \nu_x(X) + \nu_y(Y) - z = \nu(X) + \nu(Y)$. Therefore, $X + Y$ is G -linear at every generic $z \in X + Y$ and we have $\nu_z(X + Y) - z = \nu(X) + \nu(Y)$.

It follows that $(X + Y)_{lin}$ is large in $X + Y$ and the germs of $X + Y$ at all points in $(X + Y)_{lin}$ are G -equivalent. Taken together with the above, we see that $X + Y$ satisfies the assumptions of Lemma 2.12(1), hence it is G -linear and we have $\nu(X + Y) = \nu(X) + \nu(Y)$.

If X and Y are G -equivalent then $\nu(X) = \nu(Y)$, hence $\nu(X) + \nu(Y) = \nu(X)$, so $X + Y$ is G -equivalent to both X and Y .

(ii) It is easy to see that for all $z \in X \cup Y$, if $x \notin Fr(X) \cup Fr(Y)$, then we have $(X \cup Y) - z =_0 X - z$ (if $z \in X$) or $(X \cup Y) - z =_0 Y - z$ (if $z \in Y$). Because X and Y are G -linear and G -equivalent, it follows that $(X \cup Y)_{lin}$ is large in $X \cup Y$. Also, for every $x \in X$ we clearly have $X - x \subseteq_0 (X \cup Y) - x$, and similarly for $y \in Y$. We therefore can apply Lemma 2.12(1) again and conclude that $X \cup Y$ is G -linear, and G -equivalent to X and Y . \square

Lemma 2.15. *Let X be a G -linear set and Y_1, Y_2 two G -subsets of X . Then*

(i) $Y_1 + Y_2$ is a G -subset of X .

(ii) $Y_1 \cup Y_2$ is a G -subset of X .

Proof. (i) Because Y_1 and Y_2 are G -subsets of X , for all $y_1 \in Y_1$, $y_2 \in Y_2$, we have

$$(\nu_{y_1}(Y_1) + \nu_{y_2}(Y_2)) - (y_1 + y_2) = (\nu_{y_1}(Y_1) - y_1) + (\nu_{y_2}(Y_2) - y_2) \subseteq \nu(X) + \nu(X) = \nu(X).$$

Again as before, if $z = y_1 + y_2$ is a locally generic element of $Y_1 + Y_2$ then

$$\nu_z(Y_1 + Y_2) - z = \nu_{y_1}(Y_1) + \nu_{y_2}(Y_2) - z \subseteq \nu(X)$$

and hence $Y_1 + Y_2$ is G -subset of X at z . We thus have $(Y_1 + Y_2)_{slin}$ dense in $Y_1 + Y_2$. By Lemma 2.11, $Y_1 + Y_2$ is a G -subset of X .

(ii) Here we only need to note that for $z \in Y_1 \cap Y_2$, it is not true in general that the germs of Y_1 and Y_2 at z coincide. However, it is still true that $(Y_1 \cup Y_2) - z \subseteq_0 X - g$, so we can proceed as before. \square

We recall the following definition:

Definition 2.16. *Given a definable group G in a sufficiently saturated \mathcal{M} , a subgroup \mathcal{H} of G is called locally-definable, if it can be written as the directed union of definable sets $\mathcal{H} = \bigcup\{X_i : i \in I\}$, where $|I| < \kappa$.*

We say that \mathcal{H} is definably connected if the X_i 's can all be chosen to be definably connected.

Such groups were sometimes called \vee -definable groups (see [6]) or Ind-definable groups (see [4]). The dimension of a locally-definable group is taken to be $\max\{\dim X_i : i \in I\}$. Notice that if \mathcal{H} is definably connected then it is actually definably path connected in the sense that any two points can be connected by a definable path in \mathcal{H} . The following claim is easy to verify:

Claim 2.17. *If $\mathcal{H} = \bigcup\{X_i : i \in I\}$ is a locally-definable group and $g \in \mathcal{H}$ then there exists $i \in I$ such that $g \in X_i$ and*

$$\nu_g(\mathcal{H}) = g + \nu_0(\mathcal{H}) = \nu_g(X_i).$$

Lemma 2.18. *Every locally-definable subgroup \mathcal{H} of G can be written as the directed union of G -linear sets.*

Proof. Without loss of generality all X_i 's in the definition of \mathcal{H} have maximal dimension d . By Claim 2.17, for every $g \in \mathcal{H}$ there exists $i \in I$ such that $\nu_g(\mathcal{H}) = \nu_g(X_i)$. For such an i we have: $g \in (X_i)_{lin}$, the local dimension of X_i at g equals d (hence, $g \in (X_i)_{lin}^{max}$) and all $(X_i)_{lin}^{max}$ are G -linear and G -equivalent. If we let $X'_i = (X_i)_{lin}^{max}$ then we have $\mathcal{H} = \bigcup_{i \in I} X'_i$.

To see that $\{X'_i : i \in I\}$ is a directed system of sets: Given $i, j \in I$, we take $k \in I$ such that $X_i \cup X_j \subseteq X_k$ and claim that $X'_i \cup X'_j \subseteq X'_k$. Indeed, if $g \in X'_i$ then $\nu_g(X'_i) = \nu_g(\mathcal{H})$ and hence $\nu_g(X_i) = \nu_g(X_k)$. In particular, $g \in (X_k)_{lin}^{max} = X'_k$. \square

Lemma 2.19. *We assume \mathcal{M} is an ω -saturated structure. Let X be a definable G -linear set, $0 \in X$. Then the following hold.*

(1) *The group $\langle X \rangle$ generated by X is locally-definable and its germ at 0 equals to $\nu(X)$. In particular, $\dim(\langle X \rangle) = \dim X$.*

(2) *If Y a G -subset of X containing 0, then the group generated by Y is a locally-definable of dimension $\leq \dim X$, whose germ at 0 is contained in $\nu(X)$.*

Proof. (1) The group $\langle X \rangle$ generated by X is a countable increasing union of the sets $X_0 = X$, $X_1 = X - X$, $X_2 = (X - X) + (X - X), \dots$, so locally-definable. By the Lemma 2.14, each X_n is G -linear and G -equivalent to X (so in particular, has the same dimension as X). Given $g \in \langle X \rangle$, there exists $n \geq 0$ such that $g \in X_n$. Because of the G -linearity, for every $k \geq n$, we have $X_n =_g X_k$. Because of saturation, there exists a neighborhood U of g such that $U \cap \langle X \rangle = U \cap X_n$. It follows that the germ of $\langle X \rangle$ at this point equals to that of X_n and in particular, $\langle X \rangle_0 = \nu(X)$.

(2) The group $\langle Y \rangle$ generated by Y is a countable increasing union of the sets $Y_0 = Y$, $Y_1 = Y - Y$, $Y_2 = (Y - Y) + (Y - Y), \dots$. Because $-Y$ is also a G -subset of X , we can apply Lemma 2.15 and conclude that each Y_n is a G -subset of X whose germ at 0 is contained in $\nu(X)$. It follows that the dimension of $\langle Y \rangle$ is at most $\dim X$ and that the germ of $\langle Y \rangle$ at 0 is contained in that of $\langle X \rangle$. \square

We end this section with a small observation on locally-definable subgroups:

Lemma 2.20. *Let $\mathcal{H}_1, \mathcal{H}_2$ be locally-definable subgroups of G . Then $\mathcal{H}_1 + \mathcal{H}_2$ is a locally-definable group whose germ at 0 equals to the sum of the germs of \mathcal{H}_1 and \mathcal{H}_2 at 0.*

Proof. Let

$$\mathcal{H}_1 = \bigcup_{i \in I} X_i ; \quad \mathcal{H}_2 = \bigcup_{j \in J} Y_j.$$

As was pointed out above, we may assume that the X_i 's Y_j 's are all G -linear. By Lemma 2.14(1), the sets $X_i + Y_j$ are all G -linear and $\nu(X_i + Y_j) = \nu(X_i) + \nu(Y_j)$. By Lemma 2.19(1),

$$\nu_0(\mathcal{H}_1 + \mathcal{H}_2) = \nu(X_i + Y_j) = \nu(X_i) + \nu(Y_j) = \nu_0(\mathcal{H}_1) + \nu_0(\mathcal{H}_2).$$

□

3 Definable sets containing abstract subgroups

In this section \mathcal{M} will denote an ω -saturated o-minimal expansion of an ordered group.

Theorem 3.1. *Let G be a definable abelian group, $\Gamma \subseteq G$ an abstract subgroup (i.e., Γ is not necessarily definable). Let $X \subseteq G$ be a definable set containing Γ of minimal dimension d . Then there exist a definable set X' of dimension d , a definably connected locally-definable subgroup \mathcal{H} of dimension d , and $g_1, \dots, g_k \in \Gamma$ such that $\Gamma \subseteq X' \subseteq \bigcup_{i=1}^k \mathcal{H} + g_i$.*

Proof. Assume the nontrivial case $d > 0$. For every $g \in \Gamma$, the set $X \cap (g + X)$ contains Γ and, hence by minimality, has dimension d . It follows that the set $Y = \{g \in X : \dim(X \cap X + g) = d\}$ contains Γ and therefore, again by minimality, has dimension d .

We now apply Lemma 2.8 to Y, Y, X and X (for X_0, X, Y and Z , respectively), and conclude that Y_{lin} is large in Y .

Since Y contains Γ we may replace X with Y and assume from now on that X_{lin} is large in X (however, it need not be the case that X or even X_{lin} is G -linear). Moreover, we pick $X \supseteq \Gamma$ such that X_{lin} has the minimal number of definably connected components of dimension d . Note that each component has infinitely many elements of Γ (otherwise, we can replace it by finitely many points).

Claim 1 All the components of X_{lin} of dimension d are G -equivalent to each other.

Proof Indeed, if X_1, X_2 are two such components then for $g \in X_1 \cap \Gamma$, the set $\{h \in X_2 : g + h \in X\} = X_2 \cap X - g$ contains $X_2 \cap \Gamma$ and therefore has dimension d (otherwise, we could replace X_2 by a definable set of smaller dimension). For the same reason, the set of all $g \in X_1$ such that $\dim(X_2 \cap (X - g)) = d$ has dimension d . Just like in the proof of Corollary 2.8, we can apply Lemma 2.7 locally to X_1 and X_2 , and conclude that for some open U and V we have $U \cap X_1$ is G -equivalent to $V \cap X_2$. Because X_1 and X_2 are G -linear it follows that they are G -equivalent to each other.

End of Claim 1.

We therefore showed that X_{lin}^{max} is G -linear. It is left to handle $X^* = X \setminus X_{lin}^{max}$. Fix one of the components X_0 of X_{lin}^{max} .

As before, for every $g \in \Gamma \cap X^*$, the set $\{h \in X_0 : g + h \in X\}$ contains $X_0 \cap \Gamma$ and hence has dimension d . Therefore, after possibly replacing X^*

by a smaller set we may assume that for all $g \in X^*$ the set $(g + X_0) \cap X$ has dimension d .

Claim 2 X is a G -subset of X_{lin}^{max} .

Proof By abuse of notation we let $\nu(X)$ be the infinitesimal subgroup associated to the G -linear set X_{lin}^{max} . By Lemma 2.11, it is enough to see that a dense subset of X is G -linear in X_{lin}^{max} . Namely, we will show that for every locally generic $g \in X$, we have $\nu_g(X) - g \subseteq \nu(X)$. It is clearly sufficient to consider $g \in X^*$.

We fix an open set $U \subseteq G$ and $g \in X^*$ which is generic in $U \cap X$. By our assumption on X^* , there exists h generic in X_0 over g such that $g + h$ (generic) in X . As in the proof of Lemma 2.7, it follows that $\nu_g(X) + h \subseteq \nu_{g+h}(X)$, hence

$$\nu_g(X) - g \subseteq \nu_{g+h}(X) - (g + h) = \nu(X).$$

(the right-most equality follows from the fact that $g + h$ is generic in X and hence belongs to X_{lin}^{max}). **End of Claim 2.**

Let $X_{lin}^{max} = X_1 \cup \dots \cup X_r$ be the union of those components of X_{lin} of dimension d . By Lemma 2.4 each X_i is G -linear. For each X_i , pick $g_i \in \Gamma \cap X_i$, and consider the set $X'_i = X_i - g_i$. By Claim 1, the union $X' = \bigcup_{i=1}^r X'_i$ is G -linear (and $\nu(X') = \nu(X_i)$ for every i). It also contains 0 and it is definably connected. Let \mathcal{H}' be the subgroup of G generated by X' . By Lemma 2.19(1), $\dim \mathcal{H}' = \dim X' = d$. We thus have

$$X_{lin}^{max} \subseteq \bigcup_{i=1}^r \mathcal{H}' + g_i.$$

Let X^1, \dots, X^t be the definably connected components of X^* . For every such X^j , we may assume that $X^j \cap \Gamma \neq \emptyset$ (for otherwise we may omit this component), take $g^j \in \Gamma \cap X^j$, and let $X'' = \bigcup_{i=1}^t X^j - g^j$. By Claim 2, each X^j (and therefore also $X^j - g^j$) is a G -subset of X_{lin}^d . By Lemma 2.15 (2), the set X'' is also a G -subset of X_{lin}^{max} . It is also definably connected and contains 0.

Let \mathcal{H}'' be the subgroup of G generated by X'' . By Lemma 2.19(2), the germ of \mathcal{H}'' at 0 is contained in that of X' and therefore of \mathcal{H}' . We now let $\mathcal{H} = \mathcal{H}' + \mathcal{H}''$. By Lemma 2.20, we have $\nu_0(\mathcal{H}) = \nu_0(\mathcal{H}' + \mathcal{H}'') = \nu_0(\mathcal{H}')$, so in particular the dimension of \mathcal{H} is d .

Putting the above facts together we obtain:

$$\Gamma \subseteq X \subseteq \bigcup_{i=1}^k \mathcal{H} + g_i,$$

for some $g_1, \dots, g_k \in \Gamma$.

With that we end the proof of the theorem. \square

Corollary 3.2. *Let G be a definably connected abelian group. Let Γ be a divisible subgroup of the subgroup of torsion points $Tor(G)$. Let $X \subset G$ be a definable set containing Γ . Then, there is a definably connected locally-definable \mathcal{H} subgroup of G with $\dim \mathcal{H} \leq \dim X$ such that $\Gamma \subset \mathcal{H}$.*

Proof. By Theorem 3.1, taking X of minimal dimension, there is a definably connected locally-definable group \mathcal{H} subgroup of G , with $\dim \mathcal{H} = \dim X$, and $g_1, \dots, g_k \in \Gamma$ such that $\Gamma \subseteq \bigcup_{i=1}^k \mathcal{H} + g_i$. Then $\Gamma \subset \mathcal{H}$. Indeed, let $g \in \Gamma$ and let $m = \text{lcm}(o(g_1), \dots, o(g_k))$, since Γ is divisible there is $h \in \Gamma$ such that $g = mh$. For such h there is an i such that $h = h' + g_i$, for some $h' \in \mathcal{H}$. Then $g = mh = mh' \in \mathcal{H}$. \square

4 The main result: commutative case

In this section \mathcal{M} will be an ω -saturated o-minimal expansion of a real closed field and G will denote a definably compact definably connected abelian group of dimension n . Such G is divisible and hence the subgroup $Tor(G)$ and any p -Sylow of G are also divisible (the p -Sylow of G is $G_p = \bigcup_{n>0} G[p^n]$, where $G[m] = \{g \in G : mg = 0_G\}$).

By Theorem 1.1 in [3] we have $G[m] \cong (\mathbb{Z}/m\mathbb{Z})^n$ for any $m > 0$, and $\pi_1(G) \cong \mathbb{Z}^n$, where $\pi_1(G)$ is the o-minimal fundamental group of G .

Let p be a prime number and let $x_1, \dots, x_n \in G[p]$ we say that x_1, \dots, x_n are n independent p -torsion points if they are \mathbb{F}_p -independent under the isomorphism $G[p] \cong \mathbb{F}_p^n$.

Theorem 4.1. *Let X be a definable subset of G . If X contains a p -Sylow of G , then $\dim X = n$.*

Lemma 4.2. *Let p be a prime number and let $x_1, \dots, x_n \in G$ be n independent p -torsion points. For each $i = 1, \dots, n$ let τ_i be a path in G from 0_G to x_i , and let $p\tau_i$ denote the loop at 0_G , $t \mapsto p\tau_i(t)$. Then, $[p\tau_1], \dots, [p\tau_n] \in \pi_1(G)$ are \mathbb{Z} -linearly independent.*

Proof. Let $\varphi: \pi_1(G) \rightarrow G[p]$ the homomorphism $[\gamma] \mapsto \tilde{\gamma}(1)$, where $\tilde{\gamma}$ is the unique path in G starting at 0_G and such that $p\tilde{\gamma} = \gamma$ (see the proofs of propositions 2.10 and 2.11 in [3]). Suppose $[p\tau_1], \dots, [p\tau_n]$ are \mathbb{Z} -linearly dependent and let $m_1[p\tau_1] + \dots + m_n[p\tau_n] = [k_{0_G}]$ with $(m_1, \dots, m_n) = 1$, where k_{0_G} is the constant loop at 0_G . Applying φ to this equality we get $m_1x_1 + \dots + m_nx_n = 0_G$ and hence $p|(m_1, \dots, m_n)$, a contradiction. \square

Lemma 4.3. *Let $[\gamma_1], \dots, [\gamma_n] \in \pi_1(G)$ be \mathbb{Z} -linearly independent. Then,*

$$\left\{ \sum_{i=1}^n \gamma_i(t_i) : t_i \in [0, 1], 1 \leq i \leq n \right\} = G.$$

Proof. Consider the definably compact definably connected n -dimensional abelian group $\mathbb{T} = [0, 1]^n$ and the definable map

$$f: \begin{array}{ccc} \mathbb{T} & \longrightarrow & G \\ (t_1, \dots, t_n) & \mapsto & \sum_{i=1}^n \gamma_i(t_i). \end{array}$$

It suffices to prove that f is onto. Since f is continuous with respect to the manifold topology of \mathbb{T} , we can see f as a definable continuous map between definable manifolds.

Claim It suffices to prove that f induces an isomorphism $f_*: H_1(\mathbb{T}; \mathbb{Q}) \rightarrow H_1(G; \mathbb{Q})$.

Proof The map f has a degree for any orientations of \mathbb{T} and G , since both \mathbb{T} and G have dimension n (see section 4 in [3]). To prove that f is onto it suffices to prove that one (equivalently each one) of these degrees is not zero. Let ζ_G and $\zeta_{\mathbb{T}}$ be the fundamental classes of some given orientations of G and \mathbb{T} , i.e., ζ_G (respectively $\zeta_{\mathbb{T}}$) is a generator of $H_n(G) \cong \mathbb{Z}$ (resp. of $H_n(\mathbb{T}) \cong \mathbb{Z}$). The degree of f is defined by $f_*(\zeta_{\mathbb{T}}) = \deg(f)\zeta_G$. If ω_G and $\omega_{\mathbb{T}}$ are the corresponding cohomology classes by duality we have $f^*(\omega_G) = \deg(f)\omega_{\mathbb{T}}$. Hence to prove that $\deg(f) \neq 0$ suffices to prove that f induces an isomorphism of the \mathbb{Q} -vector spaces $f^*: H^n(G; \mathbb{Q}) \rightarrow H^n(\mathbb{T}; \mathbb{Q})$. Now $f^*: H^*(G; \mathbb{Q}) \rightarrow H^*(\mathbb{T}; \mathbb{Q})$ is also \mathbb{Q} -algebra homomorphism and by Theorem 1.1 in [3], both \mathbb{Q} -algebras are generated by elements of degree one. So it suffices to prove that f induces an isomorphism $f^*: H^1(G; \mathbb{Q}) \rightarrow H^1(\mathbb{T}; \mathbb{Q})$. Applying duality again we have the required result. End of Claim.

Now, let $\delta_i: [0, 1] \rightarrow \mathbb{T}: t \mapsto \delta_i(t) = (0, \dots, \overset{i}{1}, \dots, 0)$. The map f induces a map $f_*: \pi_1(\mathbb{T}) \rightarrow \pi_1(G): [\delta_i] \mapsto f_*([\delta_i]) = [\gamma_i]$, for each $i = 1, \dots, n$, which is one to one and has finite cokernel ($= \pi_1(G)/\text{Im}(f_*)$) because we have n \mathbb{Z} -linear independent $[\gamma_i]$'s. Identifying the π_1 's with the H_1 's via the Hurewicz isomorphism, we have the following exact sequence

$$0 \rightarrow H_1(\mathbb{T}) \xrightarrow{f_*} H_1(G) \rightarrow H_1(G)/\text{Im}(f_*) \rightarrow 0.$$

tensoring with \mathbb{Q} we obtain that $f_*: H_1(\mathbb{T}; \mathbb{Q}) \rightarrow H_1(G; \mathbb{Q})$ is an isomorphism. \square

Corollary 4.4. *Let \mathcal{H} be a definably connected locally-definable subgroup of G . Suppose \mathcal{H} contains $G[p]$, for some prime p . Then $\mathcal{H} = G$.*

Proof. Let x_1, \dots, x_n be n independent p -torsion points and for each $i = 1, \dots, n$ let τ_i be a path in \mathcal{H} from 0_G to x_i . Let γ_i denote the loop at 0_G defined by $t \mapsto p\tau_i(t)$, since \mathcal{H} is a group $\gamma_i([0, 1]) \subset \mathcal{H}$. By Lemma 4.2 $[\gamma_1], \dots, [\gamma_n] \in \pi_1(G)$ are \mathbb{Z} -linearly independent.

By Lemma 4.3, $S = \{\sum_{i=1}^n \gamma_i(t_i) : t_i \in [0, 1], 1 \leq i \leq n\} = G$. Again since \mathcal{H} is a group $S \subset \mathcal{H}$. \square

Proof of Theorem 4.1. Let $X \subseteq G$ be a definable set containing a p -Sylow G_p of G . Since G_p is divisible, by Corollary 3.2, there exists a definably connected locally-definable group \mathcal{H} of dimension $\leq \dim X$ containing G_p . Since G_p contains $G[p]$, we can apply Corollary 4.4 to get $\mathcal{H} = G$ and hence $\dim X = \dim G$ \square

Corollary 4.5. *Let X be a definable subset of G . If X is large in G then for any prime p and any l there is an $m > l$ and a $g \in X$ of order p^m .*

Proof. Note that this corollary is equivalent to Theorem 4.1. Indeed, for the nontrivial case suppose there is a prime p and an l such that $X \cap G_p \subset G[p^l]$. So the set $X' = X \setminus G[p^l]$ is still large in G and $X' \cap G_p = \emptyset$. But then $G \setminus X'$ contains G_p and $\dim(G \setminus X') < \dim G$, a contradiction with the theorem. \square

5 The main result: general case

We work in an ω -saturated structure \mathcal{M} which is an o-minimal expansion of a real closed field \mathcal{M} . We will use multiplicative notation for groups, and denote by p_k the map $x \mapsto x^k$. A definably connected group which is either abelian or definably compact is divisible (see [1] or [2]), hence the map p_k is onto. Recall that a *generic subset* Y of a definable abelian group G is a definable subset of G such that finitely many translates of Y cover G .

Lemma 5.1. *Let T be a definable abelian group. Suppose that for every $Y \subseteq T$ generic, $Y \cap \text{Tor}(T) \neq \emptyset$. Then, for each $Y \subseteq T$ generic there is a $k \in \mathbb{N}$ such that $p_k(Y) = T$.*

Proof. First note that for each $Y \subseteq T$ generic, there are $g_1, \dots, g_l \in \text{Tor}(T)$ such that $T = \bigcup_{i=1}^l g_i Y$. Indeed, if not, by compactness, there is $h \notin \text{Tor}(T)Y$, i.e., $hY^{-1} \cap \text{Tor}(T) = \emptyset$, but hY^{-1} is also generic, a contradiction. Now taking $k = \text{lcm}(o(g_1), \dots, o(g_l))$ we have $p_k(Y) = p_k(T) = T$ as required. \square

We recall that a maximal definably connected abelian subgroup T of a definably connected definably compact group G is called a *maximal torus* of G and that $G = \bigcup_{g \in G} T^g$ (see Theorem 6.12 in [1] or [2]).

Lemma 5.2. *Let G be a definably compact definably connected group. Let T be a maximal torus of G and $X \subseteq G$ is definable. Assume the following:*

(i) *For every $g \in G$ and for every $Y \subseteq T^g$ generic, $Y \cap \text{Tor}(T^g) \neq \emptyset$, and*

(ii) *For every $g \in G$, $X \cap T^g$ is generic in T^g .*

Then there is a $k \in \mathbb{N}$ such that $p_k(X) = G$.

Proof. Given $g \in G$, by (ii), we can apply Lemma 5.1, for $Y = X \cap T^g$, to get a $k(g) \in \mathbb{N}$ such that the map $y \mapsto y^{k(g)}$ sends $X \cap T^g$ onto T^g (note that by divisibility of T^g any multiple of $k(g)$ has the same property). Then, by compactness, there is $k \in \mathbb{N}$ such that for every $g \in G$, the map $y \mapsto y^k$ sends $X \cap T^g$ onto T^g . To finish the proof suffices to make use of the equality $G = \bigcup_{g \in G} T^g$. \square

For the next lemma we will use the notions of *compact domination* and *very good reduction* introduced in [4]. Recall that a definably simple group has very good reduction and that a definably compact group which has very good reduction has also compact domination (see Theorem 10.7 in [4]).

Lemma 5.3. *Let G be a definably compact definably connected group. Let $X \subseteq G$ be a definable set such that $\text{Tor}(G) \subseteq X$. If G has very good reduction then there is $k \in \mathbb{N}$ such that $p_k(X) = G$.*

Moreover, if N is a finite (central) normal subgroup of G and G/N has very good reduction then there is $k \in \mathbb{N}$ such that $p_k(X) = G$.

Proof. The result can probably be read off standard Lie theory. However, since we could not find a reference we give a complete proof.

Without loss of generality we may assume that $G = G(\mathcal{M})$ is defined over the reals. By our assumptions on G , $G(\mathbb{R})$ is a compact connected Lie group. Let T_0 be a standard maximal torus of $G(\mathbb{R})$, hence T_0 is also a Lie group and hence definable over the reals. Let $T = T_0(\mathcal{M})$. So T is still a maximal torus of G and it has very good reduction and hence it is compactly dominated. The same happens for every conjugate T^g of T , for $g \in G$. Since all T^g 's are also abelian, we can apply Proposition 10.6 in [4], to get that for every $g \in G$, every generic subset of T^g has a torsion point, so condition (i) of Lemma 5.2 is satisfied.

Because $Tor(G) \subseteq X$, the set $T^g \setminus X$ has no torsion and hence, by (i), it is non-generic. It follows that $T^g \cap X$ is generic and therefore (ii) holds. Now we can apply Lemma 5.2 to get the required result.

For the moreover clause, let $\pi : G \rightarrow G/N$ be the projection map. Because N is finite, all torsion elements of G/N are in $\pi(X)$, hence by what we have just proved, there is k such that $p_k(\pi(X)) = G/N$. If we take $k' = k \cdot |N|$ then $p_{k'}(X) = G$ (because N is central and G divisible). \square

Notice, in the setting of the above lemma, that if $p_k(X) = G$ then in particular, $\dim X = \dim G$ (since the image of X under a definable map cannot increase).

We can now prove the main result:

Theorem 5.4. *Let G be a definably compact group and assume that $X \subseteq G$ is a definable set containing all torsion points of G . Then $\dim X = \dim G$.*

Proof. By 4.1 in [5] and 5.4 in [8], after modding out $G_1 = G/Z(G)$ by its finite center, the group we obtain is a direct product of simple groups. In particular, it has very good reduction (see the proof of 5.1 in [7]) and hence the “moreover” clause of Lemma 5.3 holds for $G/Z(G)$. We therefore have:

(i) For every definable $X_1 \subseteq Z(G)$, if $Tor(Z(G)) \subseteq X_1$ then $\dim X_1 = \dim Z(G)$ (by Theorem 4.1 this is true for subsets of $Z(G)^0$, but then it clearly follows for subsets of $Z(G)$ as well).

(ii) For every definable $X_1 \subseteq G$, if $Tor(G/Z(G)) \subseteq X_1/Z(G)$ then $\dim(X_1/Z(G)) = \dim(G/Z(G))$. (by Lemma 5.3 and our previous observation).

We now proceed as follows, with $X \subseteq G$ a definable set containing $Tor(G)$. Given $g \in Tor(G)$ and $h \in Tor(Z(G))$, we have $gh \in Tor(G) \subseteq X$. Hence, for every such g , we have

$$X_g = \{h \in Z(G) : gh \in X\} \supseteq Tor(Z(G)),$$

so, by (i), $\dim X_g = \dim Z(G)$. Since $gX_g = (gZ(G)) \cap X$, we have, for every $g \in Tor(G)$,

$$\dim gZ(G) \cap X = \dim Z(G).$$

Let

$$X_1 = \{g \in G : \dim(gZ(G) \cap X) = \dim Z(G)\}.$$

It follows from the above that $Tor(G) \subseteq X_1$, and therefore $Tor(G/Z(G)) \subseteq X_1/Z(G)$. (Indeed, if $hZ(G) \in Tor(G/Z(G))$ then for some $n \in \mathbb{N}$, $h^n \in Z(G)$, and therefore for some $k \geq n$, $h^k \in Z(G)^0$. Because $Z(G)^0$ is divisible there exists $h_1 \in Z(G)$, with $h_1^k = h^k$. Because h_1 is central,

$(hh_1^{-1})^k = e$, hence $hh_1^{-1} \in X_1$ and therefore $hZ(G) \in X_1/Z(G)$. By (ii), $\dim(X_1/Z(G)) = \dim(G/Z(G))$.

To finish the proof, consider the set $Y = (X_1Z(G)) \cap X$. The cosets of $Z(G)$ partition Y into equivalence classes, each of dimension $\dim Z(G)$ (by definition of X_1). Since every $Z(G)$ -coset of an element in X_1 intersects X nontrivially, we have

$$\dim(Y/Z(G)) = \dim(X_1/Z(G)) = \dim(G/Z(G)).$$

Summarizing, we have

$$\dim Y \geq \dim Z(G) + \dim(G/Z(G)) = \dim G,$$

and hence (since $Y \subseteq X$) $\dim X = \dim G$. □

References

- [1] A.Berarducci, Zero-groups and maximal tori in **Logic Colloquium'04**, A. Andretta *et al.* (eds.),ASL Lecture Notes in Logic, Vol. 29 (2006) 33-45
- [2] M.Edmundo, A remark on divisibility of definable groups, *Math. Logic Quart.*51 (2005) 639-641.
- [3] M.Edmundo and M.Otero, Definably compact abelian groups, *J. Math. Logic* 4 (2004) 163-180.
- [4] E.Hrushovski, Y.Peterzil and A.Pillay, Groups, measures and NIP, *J. Am. Math.Soc.*, 34pp. to appear.
- [5] Y.Peterzil, A.Pillay and S.Starchenko, Definably simple groups in \mathcal{o} -minimal structures, *Trans. Am. Math. Soc.* 352 (2000) 4397-4419.
- [6] Y.Peterzil, A.Pillay and S.Starchenko, Simple algebraic and semialgebraic groups over real closed fields, *Trans.AMS* 352 (2000) 4421-4450.
- [7] Y.Peterzil, A.Pillay and S.Starchenko, Linear groups definable in \mathcal{o} -minimal structures, *J. Algebra* 247 (2002) 1-23.
- [8] Y.Peterzil and S.Starchenko, Definable homomorphisms of abelian groups in \mathcal{o} -minimal structures, *Ann. Pure Appl. Logic* 101 (2000) 1-27.