

BOUNDING THE CONSISTENCY STRENGTH OF A FIVE ELEMENT LINEAR BASIS

BY

BERNHARD KÖNIG*

*Department of Mathematics, University of Toronto
Toronto, Ontario M5S 2E4, Canada
e-mail: bkoenig@math.toronto.edu*

AND

PAUL LARSON**, †, ††, ‡

*Department of Mathematics and Statistics, Miami University
Oxford, Ohio 45056, USA
e-mail: larsonpb@muohio.edu*

AND

JUSTIN TATCH MOORE††

*Department of Mathematics, Cornell University
Ithaca, NY 14853, USA
e-mail: justin@math.cornell.edu*

AND

BOBAN VELIČKOVIĆ‡

*Equipe de Logique Mathématique, Université de Paris 7 Denis Diderot
75251 Paris, France
e-mail: boban@logique.jussieu.fr*

* The first author acknowledges a fellowship granted by the French ministry of research.

** The research of the second author was partially supported by the *Centre de Rercerca Matemàtica* of the *Universitat Autònoma de Barcelona*, and by NSF Grant DMS-0401603.

† The second author would also like to thank the third and fourth authors for bringing him to Boise and Paris respectively for further discussions.

†† The third author was supported by NSF grants DMS-0401893 and DMS-0200671.

‡ The second and fourth authors would like to thank CIRM in Luminy for hosting them during a *petit group de travaille*, and to thank the others participants, Ralf Schindler and Ernest Schimmerling, for discussions on this topic.

Received January 10, 2006 and in revised form July 27, 2006

ABSTRACT

In [13] it was demonstrated that the Proper Forcing Axiom implies that there is a five element basis for the class of uncountable linear orders. The assumptions needed in the proof have consistency strength of at least infinitely many Woodin cardinals. In this paper we reduce the upper bound on the consistency strength of such a basis to something less than a Mahlo cardinal, a hypothesis which can hold in the constructible universe L .

A crucial notion in the proof is the **saturation of an Aronszajn tree**, a statement which may be of broader interest. We show that if all Aronszajn trees are saturated and $\text{PFA}(\omega_1)$ holds, then there is a five element basis for the uncountable linear orders. We show that $\text{PFA}(\omega_2)$ implies that all Aronszajn trees are saturated and that it is consistent to have $\text{PFA}(\omega_1)$ plus every Aronszajn tree is saturated relative to the consistency of a reflecting Mahlo cardinal. Finally we show that a hypothesis weaker than the existence of a Mahlo cardinal is sufficient to force the existence of a five element basis for the uncountable linear orders.

1. Introduction

In [13] it was shown that the Proper Forcing Axiom (PFA) implies that the class of uncountable linear orders has a five element basis, i.e., that there is a list of five uncountable linear orders such that every uncountable linear order contains an isomorphic copy of one of them. This basis consists of X , ω_1 , ω_1^* , C , and C^* where X is any suborder of the reals of cardinality ω_1 and C is any Countryman line. In fact any five element basis for the uncountable linear orders must have this form.

Recall that a **Countryman line** is a linear order whose square is the union of countably many non-decreasing relations. These linear orders are necessarily Aronszajn. Their existence was proved by Shelah who conjectured that every Aronszajn line consistently contains a Countryman suborder [14]. In [1, p. 79] it is stated that, assuming PFA, every Aronszajn line contains a Countryman suborder if and only if the Coherent Tree Axiom (CTA) holds:

There is an Aronszajn tree T such that for every $K \subseteq T$ there is an uncountable antichain $X \subseteq T$ such that $\wedge(X)$ is either contained in or disjoint from K .

Here $\wedge(X)$ denotes the set of all pairwise meets of elements of X . Over time, this conjecture developed in the folklore and at some point it was known to

be equivalent, modulo PFA, to the assertion that the above list forms a basis for the class of uncountable linear orders.¹ Moreover, this reduction does not require any of the large cardinal strength of PFA. The reader is referred to the final section of [18, §4.4] for proofs of the above assertions.

In [13] it is shown that PFA implies CTA. In fact, the conjunction of the Bounded Proper Forcing Axiom (PFA(ω_1)) [8] and the Mapping Reflection Principle (MRP) [12] suffices. Since the consistency of MRP requires considerably large cardinal assumptions² and since it is required only in Lemma 5.29 of [13], it is natural to ask what large cardinals, if any, are necessary for CTA. Here we reduce the consistency strength needed to something less than a Mahlo cardinal,³ a hypothesis which can hold in the constructible universe L .

The main goal of this paper is to prove this Key Lemma from a greatly reduced hypothesis. A central notion in our analysis is that of the **saturation** of an Aronszajn tree T — whenever \mathcal{A} is a collection of uncountable downward closed subsets of T which have pairwise countable intersection, then \mathcal{A} has cardinality at most ω_1 . When possible, this notion will be considered separately, as it seems that this could be relevant in other contexts.

Central to the proof of the main result in [13] is the notion of **rejection**. This will be defined after recalling some preliminary definitions from [13]. For the moment, fix an Aronszajn tree $T \subseteq 2^{<\omega_1}$ which is coherent, special and closed under finite modifications,⁴ and let K be a subset of T .

Definition 1.1: If P is a countable elementary submodel of $H(\omega_2)$ containing T , let $\mathcal{I}_P(T)$ be the collection of all $I \subseteq \omega_1$ such that for some uncountable $Z \subseteq T$ in P and some t in the downwards closure of Z having height $P \cap \omega_1$, I is disjoint from

$$\Delta(Z, t) = \{\Delta(s, t) : s \in Z\}.$$

It is always the case that $\mathcal{I}_P(T)$ is closed under finite unions and subsets. The former property uses the properties of T and is non-trivial; the argument

¹ It seems that the speculation of the consistent existence of a finite basis for the uncountable linear orders first appeared in print in [3], although, its equivalence to Shelah's conjecture was unknown at that point.

² The current upper bound is a supercompact cardinal [12].

³ A regular cardinal κ is **Mahlo**, if the set of regular cardinals less than κ is stationary in κ .

⁴ The tree $T(\mathfrak{q}_3)$ of [18] is such an example.

is similar to the proof of [15, Lemma 4.1]. Similarly, $\mathcal{S}_P(T)$ remains the same if one takes t to be a fixed member of $T_{P \cap \omega_1}$ instead of letting t vary.

Definition 1.2: If X is a finite subset of T , $K(X)$ is the set of all γ which are less than the heights of all elements of X and satisfy $s \upharpoonright \gamma \in K$ for all s in X .

Definition 1.3: If X is a finite subset of T and P is a countable elementary submodel of $H(\omega_2)$, then P **rejects** X if $K(X \setminus P)$ is in $\mathcal{S}_P(T)$.

It is shown in [13] that the following lemma (Lemma 5.29 of [13]), taken in conjunction with $\text{PFA}(\omega_1)$, is sufficient to prove the existence of an uncountable antichain $X \subseteq T$ such that $\wedge(X)$ is contained in or disjoint from K .

KEY LEMMA 1.4 [13]: (MRP) *If M is a countable elementary submodel of $H(2^{2^{\omega_1}+})$ which contains T and K and X is a finite subset of T , then there exists a closed unbounded set E of countable elementary submodels of $H(\omega_2)$ such that E is in M and either every element of $E \cap M$ rejects X or no element of $E \cap M$ rejects X .*

We will begin by defining a combinatorial statement φ in Section 2 and showing that this statement implies the Key Lemma. The statement φ is a strengthening of **Aronszajn tree saturation** — the assertion that every Aronszajn tree is saturated. Moreover, φ is shown to be equivalent to Aronszajn tree saturation in the presence of $\text{PFA}(\omega_1)$. In Section 3 we will demonstrate that $\text{PFA}(\omega_2)$ implies φ . This reduces the upper bound on the consistency strength of Shelah's conjecture to something less than the existence of 0^\sharp but greater than a weakly compact cardinal. Section 4 further refines the argument to show that an instance of φ can be forced by a proper forcing without a need for large cardinal assumptions. This is then implemented in Section 5 to further optimize the upper bound on the consistency strength of a five element basis for the uncountable linear orders to something less than the existence of a Mahlo cardinal.

The notation and terminology used in this paper is fairly standard. All ordinals are von Neumann ordinals; they are the sets of their predecessors. The cardinal \beth_α is defined recursively so that $\beth_0 = \omega$, $\beth_{\alpha+1} = 2^{\beth_\alpha}$, and $\beth_\delta = \sup_{\alpha < \delta} \beth_\alpha$ for limit δ . The reader is referred to [9] as a general reference for Set Theory. In this paper, **Aronszajn tree** or **A-tree** will mean an uncountable tree in which all levels and chains are countable. A **subtree** of an A-tree T is

an uncountable downward closed subset of T . The reader is referred to [10] or [17] for further reading on bounded fragments of PFA.

2. Aronszajn tree saturation

Recall the notion of the **saturation of** $\mathcal{P}(\omega_1)/\text{NS}$:

Any collection of stationary sets, which have pairwise non-stationary intersection, has cardinality at most ω_1 .

Now consider the following statement $\psi_{\text{NS}}(\mathcal{A})$ for a collection \mathcal{A} of subsets of ω_1 :

There is a club $E \subseteq \omega_1$ and a sequence $\langle A_\xi : \xi < \omega_1 \rangle$ of elements of \mathcal{A} such that for all δ in E , there is a $\xi < \delta$ with δ in A_ξ .

The assertion ψ_{NS} that $\psi_{\text{NS}}(\mathcal{A})$ holds for every predense set $\mathcal{A} \subseteq \mathcal{P}(\omega_1)/\text{NS}$ is in fact equivalent to the saturation of $\mathcal{P}(\omega_1)/\text{NS}$. This was used to prove that Martin's Maximum implies that $\mathcal{P}(\omega_1)/\text{NS}$ is saturated, [7]. The significance of $\psi_{\text{NS}}(\mathcal{A})$, from our point of view, is that it implies that \mathcal{A} is predense and is Σ_1 in complexity and, therefore, upwards absolute.

In this section we will be interested in analogous assertions about subtrees of an A-tree.⁵

Definition 2.1: An A-tree T is **saturated** if whenever \mathcal{A} is a collection of subtrees T which have pairwise countable intersection, \mathcal{A} has cardinality at most ω_1 .

This statement follows from the stronger assertion shown by Baumgartner to hold after Levy collapsing an inaccessible cardinal to ω_2 [4].

For every A-tree T , there is a collection \mathcal{B} of subtrees of T such that \mathcal{B} has cardinality ω_1 and every subtree of T contains an element of \mathcal{B} .

Unlike the case $\mathcal{P}(\omega_1)/\text{NS}$, the maximality of an ω_1 -sized antichain of subtrees can be shown not to be upwards absolute.⁶ However, this leaves open the

⁵ Actually, all statements about the saturation of an A-tree make sense in the broader context of all ω_1 -trees. We will not need this generality and in fact the saturation of all A-trees implies the more general case by Todorćević's construction presented in Section 2 of [4].

⁶ This can be derived from the arguments in [16, §8] and the construction in [4, §2]. An explicit argument for this can be found in [11].

question of how to obtain the consistency of A-tree saturation in the presence of a forcing axiom. It is not difficult to show that both Chang's Conjecture and the saturation of $\mathcal{P}(\omega_1)/\text{NS}$ each imply that all A-trees are saturated. Therefore, Martin's Maximum implies that all A-trees are saturated. We will pursue a different proof which is weaker in terms of consistency strength and somewhat different in character.

If \mathcal{F} is a collection of subtrees of T , then \mathcal{F}^\perp is the collection of all subtrees B of T such that for every A in \mathcal{F} , $A \cap B$ is countable. If \mathcal{F}^\perp is empty, then \mathcal{F} is said to be **predense**. For \mathcal{F} , a collection of subtrees of an A-tree T , we define the following assertions:

- $\psi(\mathcal{F})$ There is a closed unbounded set $E \subseteq \omega_1$ and a continuous chain $\langle N_\nu : \nu \in E \rangle$ of countable subsets of \mathcal{F} such that for every ν in E and t in T_ν there is a $\nu_t < \nu$ such that if $\xi \in (\nu_t, \nu) \cap E$, then there is $A \in \mathcal{F} \cap N_\xi$ such that $t \upharpoonright \xi$ is in A .
- $\varphi(\mathcal{F})$ There is a closed unbounded set $E \subseteq \omega_1$ and a continuous chain $\langle N_\nu : \nu \in E \rangle$ of countable subsets of $\mathcal{F} \cup \mathcal{F}^\perp$ such that for every ν in E and t in T_ν either
- (1) there is a $\nu_t < \nu$ such that if $\xi \in (\nu_t, \nu) \cap E$, then there is $A \in \mathcal{F} \cap N_\xi$ such that $t \upharpoonright \xi$ is in A , or
 - (2) there is a B in $\mathcal{F}^\perp \cap N_\nu$ such that t is in B .

The following proposition, together with Lemma 4.3 below, captures the important properties of $\psi(\mathcal{F})$.

LEMMA 2.2: *Let \mathcal{F} be a fixed family of trees. $\psi(\mathcal{F})$ is a Σ_1 -formula with parameters \mathcal{F} , T , and ω_1 which implies that \mathcal{F} is predense.*

Proof. Let ψ_0 be the conjunction of the following formulas:

$$\begin{aligned} & \forall \alpha \in E (\alpha \in \omega_1) \\ & \forall \alpha \in \omega_1 \exists \beta \in E (\alpha \in \beta) \\ & \forall \alpha \in \omega_1 (\neg(\alpha \in E) \rightarrow \exists \beta \in \alpha \forall \gamma \in \alpha ((\gamma \in E) \rightarrow (\gamma \in \beta))). \end{aligned}$$

Clearly, ψ_0 asserts that E is a closed unbounded subset of ω_1 . Similarly, the assertion $\vec{N} = \langle N_\nu : \nu \in E \rangle$ is a continuous chain of countable subsets of \mathcal{F} is a Σ_0 -formula ψ_1 with parameters E and \mathcal{F} . Finally let ψ_2 be the Σ_0 -formula asserting that for every ν in E and t in T_ν there is a $\nu_t < \nu$ such that if $\xi \in (\nu_t, \nu) \cap E$, then there is $A \in \mathcal{F} \cap N_\xi$ such that $t \upharpoonright \xi$ is in A . Then $\psi(\mathcal{F})$

is the Σ_1 -formula

$$\exists E \exists \vec{N} \psi_0 \wedge \psi_1 \wedge \psi_2.$$

In order to see that $\psi(\mathcal{F})$ implies that \mathcal{F} is predense, suppose that S is a subtree of T and that $\langle N_\nu : \nu \in E \rangle$ witnesses $\psi(\mathcal{F})$. Let M be a countable elementary submodel of $H(2^{\omega_1+})$ which contains T , S and $\langle N_\nu : \nu \in E \rangle$ as elements. Set $\delta = M \cap \omega_1$ and select a t in S of height δ . By choice of $\langle N_\nu : \nu \in E \rangle$, there is $\delta_t < \delta$ such that if $\xi \in (\delta_t, \delta) \cap E$, then there is an A in $\mathcal{F} \cap N_\xi$ with $t \upharpoonright \xi$ in A . Let \bar{N} be a countable elementary submodel of $H(\omega_2)$ such that \bar{N} is in M and T , S , $\langle N_\nu : \nu \in E \rangle$, and δ_t are in \bar{N} . By the continuity assumption on $\langle N_\nu : \nu \in E \rangle$ and elementarity of \bar{N} , $\nu = \bar{N} \cap \omega_1$ is in E and

$$N_\nu \cap \mathcal{F} = \bigcup_{\xi \in \nu \cap E} N_\xi \cap \mathcal{F} = \bar{N} \cap \mathcal{F}.$$

Hence, there is an A in $\mathcal{F} \cap \bar{N}$ such that $t \upharpoonright \nu$ is in A . Since $t \upharpoonright \nu$ is in $S \cap A$ but not in \bar{N} , the elementarity of \bar{N} implies that $S \cap A$ must be uncountable, finishing the proof. ■

While $\varphi(\mathcal{F})$ and $\psi(\mathcal{F})$ are equivalent if \mathcal{F} is predense, $\varphi(\mathcal{F})$ is, in general, not a Σ_1 -formula in \mathcal{F} and T . Let φ be the assertion that whenever T is an A-tree and \mathcal{F} is a family of subtrees T , $\varphi(\mathcal{F})$ holds and let ψ be the analogous assertion but with quantification only over \mathcal{F} which are predense. As noted, φ implies ψ . Also, if \mathcal{A} is a predense family of subtrees of T which have pairwise countable intersects and $\langle N_\nu : \nu \in E \rangle$ witnesses $\psi(\mathcal{A})$, then $\bigcup_{\nu \in E} N_\nu = \mathcal{A}$ and, in particular, \mathcal{A} has size at most ω_1 . Hence both ϕ and ψ imply A-tree saturation.

We will now show the relevance of φ to our main goal. Before proceeding, it will be useful to reformulate the notion of rejection presented in the introduction.

Definition 2.3: Let $T^{[n]}$ denote the collection of all elements τ of T^n such that every coordinate of τ has the same height and, when considered as a sequence of elements of T , the coordinates of τ are non-decreasing in the lexicographical order on T . $T^{[n]}$ will be considered as a tree with the coordinate-wise partial order induced by T .

Remark 2.4: Intuitively, elements of $T^{[n]}$ are n -element subsets of T . In order to ensure that $T^{[n]}$ is closed under taking restrictions, it is necessary to allow for

n -element sets to have repetitions and the above definition is a formal means to accommodate this. We will abuse notation and identify elements of $T^{[n]}$ which have distinct coordinates with the set of their coordinates. In our arguments, only the range of these sequences will be relevant.

Definition 2.5: Let $n < \omega$ be fixed. For any uncountable set $Z \subseteq T$, let R_Z be downward closure of the set of elements Y of $T^{[n]}$ such that $\Delta(Z, t) \cap K(Y) = \emptyset$ for some element t of the downward closure of Z with $\text{height}(t) = \text{height}(Y)$. Let \mathcal{R}_n denote the collection of all R_Z as Z ranges over the uncountable subsets of T .

LEMMA 2.6: Suppose that P is a countable elementary submodel of $H(\omega_2)$ which has T as a member. For any $Y \in T^{[n]}$ with $\text{height}(Y) \geq P \cap \omega_1$, P rejects Y if and only if $Y \upharpoonright (P \cap \omega_1)$ is in R_Z for some $Z \in P$.

Proof. Suppose first that P rejects Y . Then there exist $Z \subseteq T$ in P and $t \in (Z \cap T_{P \cap \omega_1}) \setminus P$ such that $\Delta(Z, t) \cap K(Y) = \emptyset$. Since $K(Y) \cap (P \cap \omega_1) = K(Y \upharpoonright (P \cap \omega_1))$, it follows that $Y \upharpoonright (P \cap \omega_1)$ is in R_Z .

In the other direction, if $Y \upharpoonright (P \cap \omega_1)$ is in R_Z for some $Z \in P$, then there exist $Y' \geq Y \upharpoonright (P \cap \omega_1)$ in R_Z and $t \in Z \cap T$ such that $\text{height}(t) \geq \text{height}(Y')$ and $\Delta(Z, t) \cap K(Y') = \emptyset$. Then $K(Y') \cap (P \cap \omega_1) = K(Y \upharpoonright (P \cap \omega_1))$ and $\Delta(Z, t) \cap (P \cap \omega_1) = \Delta(Z, t \upharpoonright (P \cap \omega_1))$. Hence $\Delta(Z, t \upharpoonright (P \cap \omega_1)) \cap K(Y) = \emptyset$, which means that P rejects Y . ■

LEMMA 2.7: Suppose that T is a coherent A -tree which is closed under finite changes. If $\varphi(\mathcal{R}_n)$ holds for every $n < \omega$, then the Key Lemma holds for T .

Proof. Assume the hypothesis of the lemma and let M and X be given as in the statement of the Key Lemma. Without loss of generality, we may assume that X is in $T_{M \cap \omega_1}^{[n]}$ for some $n < \omega$. Note that \mathcal{R}_n is Σ_1 -definable using parameters for T and K and, therefore, there is an $\vec{N} = \langle N_\nu : \nu \in E \rangle$ in M which witnesses $\varphi(\mathcal{R}_n)$. Either there is a $\nu_X < M \cap \omega_1$ such that for each $\xi \in E \cap (\nu_X, M \cap \omega_1)$ there is an $R \in N_\xi \cap \mathcal{F}$ with $X \upharpoonright \xi \in R$ or there is a B in $\mathcal{R}_n^\perp \cap N_{M \cap \omega_1}$ with $X \in B$.

In the first case, let E be the set of countable P elementary submodels of $H(\omega_2)$ which satisfy $\vec{N} \in P$ and $P \cap \omega_1 > \nu_X$. Then every member P of E contains $N_{P \cap \omega_1}$ and thus contains an $R \in \mathcal{R}_n$ with $X \upharpoonright (P \cap \omega_1)$ in R . By Claim 2.6, P rejects X .

In the second case, let E be the set of all P countable elementary submodels of $H(\omega_2)$ with K, T , and B in P . If P in $E \cap M$ were to reject X , there would be an R in $\mathcal{R}_n \cap P$ with $X \upharpoonright (P \cap \omega_1) \in R$. It would follow that $X \upharpoonright (P \cap \omega_1)$ is in $B \cap R$, which by elementarity of P would imply that $B \cap R$ is uncountable which is contrary to B being in \mathcal{R}_n^\perp . Hence no element of $E \cap M$ rejects X . ■

3. PFA(ω_2) implies φ

In this section we will show that PFA(ω_2) implies φ . If λ is a cardinal, then PFA(λ) is the fragment of PFA in which only antichains of size at most λ are considered, [8]. We will use the following reformulation which is due to Miyamoto, [10]:

PFA(λ): For every A in $H(\lambda^+)$ and Σ_0 -formula ϕ , if some proper partial order forces $\exists X \phi(X, A)$, then there is a stationary set of N in $[H(\lambda^+)]^{\omega_1}$ such that A is in N and $H(\omega_2)$ satisfies $\exists X \phi(X, \pi_N(A))$, where π_N is the transitive collapse of N .

In [10] it is also shown that PFA(ω_2) is equiconsistent with the existence of a cardinal κ which is $H(\kappa^+)$ -reflecting. Such cardinals are larger than weakly compact cardinals but still relativize to L and hence do not imply the existence of 0^\sharp . In this section we show that PFA(ω_2) implies that every A-tree is saturated. First, we recall some definitions from [12].

Definition 3.1: Let θ be a regular cardinal, let X be uncountable and let M be a countable subset of $H(\theta)$ such that $[X]^\omega$ is in M . A subset Σ of $[X]^\omega$ is **M -stationary** if and only if for all E in M such that $E \subseteq [X]^\omega$ is club, $\Sigma \cap E \cap M$ is non-empty.

The **Ellentuck topology** on $[X]^\omega$ is obtained by declaring a set **open** if and only if it is the union of sets of the form

$$[x, N] = \{Y \in [X]^\omega : x \subseteq Y \subseteq N\}$$

where $N \in [X]^\omega$ and $x \subseteq N$ is finite. When we say ‘open’ in this paper we refer to this topology.

Definition 3.2: A set mapping Σ is **open stationary** if and only if there is an uncountable set $X = X_\Sigma$ and a regular cardinal $\theta = \theta_\Sigma$ such that $[X]^\omega \in H(\theta)$,

$\text{dom}(\Sigma)$ is a club in $[H(\theta)]^\omega$ and $\Sigma(M) \subseteq [X]^\omega$ is open and M -stationary, for every M in the domain of Σ .

Definition 3.3: Suppose Σ is an open stationary set mapping. We say that $\langle N_\xi : \xi < \omega_1 \rangle$ is a **reflecting sequence for Σ** , if it is a continuous \in -chain contained in the domain of Σ such that for all limit $\nu < \omega_1$, there is a $\nu_0 < \nu$ such that if $\nu_0 < \xi < \nu$, then $N_\xi \cap X$ is in $\Sigma(N_\nu)$.

The **Mapping Reflection Principle** (MRP) is the assertion that every open stationary set mapping which is defined on a club admits a reflecting sequence. In [12] it was shown that MRP follows from PFA, by demonstrating the following theorem which will be useful to us here.

THEOREM 3.4: [12] *If Σ is an open stationary set mapping defined on a club, then there is a proper forcing which adds a reflecting sequence for Σ .*

LEMMA 3.5: *Suppose that T is a saturated A -tree. Then for every family \mathcal{F} of subtrees of T there is a subfamily \mathcal{F}' of \mathcal{F} of cardinality at most ω_1 such $\mathcal{F}^\perp = (\mathcal{F}')^\perp$.*

Proof. Supposing that the lemma is false, for each $\alpha < \omega_2$, recursively choose subtrees F_α, R_α of T such that each $F_\alpha \in \mathcal{F}$, each $R_\alpha \cap F_\alpha$ is uncountable, and $R_\alpha \cap F_\beta$ is countable for each $\beta < \alpha$. Then the trees $F_\alpha \cap R_\alpha$ form an antichain of cardinality ω_2 . ■

LEMMA 3.6: *Let κ be a cardinal greater than or equal to $(2^{\omega_1})^+$. Suppose that T is an A -tree, \mathcal{F} is a collection of subtrees of T and M is a countable subset of $H(\kappa)$ which has T and \mathcal{F} as members and satisfies all axioms of ZFC except the power set axiom. If t is an element of T of height $M \cap \omega_1$ and*

$$\{P \in [H(\omega_2)]^\omega : \exists A \in \mathcal{F} \cap P (t \upharpoonright (P \cap \omega_1) \in A)\}$$

is not M -stationary, then there is an S in $M \cap \mathcal{F}^\perp$ which contains t .

Proof. Let $\delta = M \cap \omega_1$. Let $E \in M$ be a club of countable elementary submodels of $H(\omega_2)$ such that $E \cap M$ is disjoint from

$$\{P \in [H(\omega_2)]^\omega : \exists A \in \mathcal{F} \cap P (t \upharpoonright (P \cap \omega_1) \in A)\}.$$

Let S be the set of all $s \in T$ such that there exist no P in E with $P \cap \omega_1 < \text{height}(s)$ and A in $\mathcal{F} \cap P$ such that $s \upharpoonright (P \cap \omega_1)$ is in A . We claim that $t \in S$. Otherwise, there would exist a P in E with $P \cap \omega_1 < \delta$ and an A in $\mathcal{F} \cap P$

such that $t \upharpoonright (P \cap \omega_1)$ is in A . Letting $\gamma = P \cap \omega_1$, this is a statement about $t \upharpoonright \gamma$, which is an element of M , so by the elementarity of M there would exist a P in $E \cap M$ with $P \cap \omega_1 = \gamma$ and an A in $\mathcal{F} \cap P$ such that $t \upharpoonright \gamma$ is in A , contradicting our choice of E . Therefore t is in S .

Clearly S is downwards closed and it is uncountable since it is an element of M but not a subset of M . We are finished once we show that $S \cap A$ is countable for every element A of \mathcal{F} . Suppose not. Since S is in M , by elementarity there must be such an A in $M \cap \mathcal{F}$. Let P be an element of E which contains both A and S . Since $A \cap S$ is uncountable and downwards closed, there must be an s in $A \cap S$ of height $(P \cap \omega_1) + 1$. But this contradicts the definition of S . ■

THEOREM 3.7: $\text{PFA}(\omega_2)$ implies φ .

Proof. In [12], it is shown that $\text{PFA}(\omega_1)$ implies that $2^{\omega_1} = \omega_2$ and hence $\text{PFA}(\omega_2)$ is equivalent to $\text{PFA}(2^{\omega_1})$. Let

$$T = \{\tau(\alpha, i) : \alpha < \omega_1 \text{ and } i < \omega\}$$

be such that $\tau(\alpha, i)$ is of height α for every α and i . Define $\Sigma_{\mathcal{F}}^i$ as follows. The domain of $\Sigma_{\mathcal{F}}^i$ is the set of all countable subsets M of $H(2^{\omega_1+})$ such that $M \cap H(\omega_2)$ is an elementary submodel of $H(\omega_2)$, \mathcal{F} and \mathcal{F}^\perp are in M , and M satisfies that $\mathcal{F} \cup \mathcal{F}^\perp$ is predense. If M is in the domain of $\Sigma_{\mathcal{F}}^i$ and there is no S in $M \cap \mathcal{F}^\perp$ with $\tau(M \cap \omega_1, i)$ in S , then let $\Sigma_{\mathcal{F}}^i(M)$ be the collection of all $P \in [H(\omega_2)]^\omega$ such that either $P \cap \omega_1$ is not an ordinal or else there is an A in $P \cap \mathcal{F}$ such that $\tau(M \cap \omega_1, i) \upharpoonright (P \cap \omega_1)$ is in A . If there is an S in $M \cap \mathcal{F}^\perp$ with $\tau(M \cap \omega_1, i)$ in S , put $\Sigma_{\mathcal{F}}^i(M)$ to be all of $[H(\omega_2)]^\omega$. Lemma 3.6 implies that $\Sigma_{\mathcal{F}}^i$ is open and M -stationary.

We will now argue that $\text{PFA}(2^{\omega_1})$ implies that each $\Sigma_{\mathcal{F}}^i$ admits a reflecting sequence. By the proof of Theorem 3.1 of [12], there is a proper forcing which introduces a reflecting sequence for each $\Sigma_{\mathcal{F}}^i$. Following the proof of Lemma 2.2, let $\phi_i(\vec{N}, \mathcal{F}, \mathcal{F}^\perp, \tau)$ be a Σ_1 -formula asserting that \vec{N} is a reflecting sequence for $\Sigma_{\mathcal{F}}^i$. While $\Sigma_{\mathcal{F}}^i$ is not an element of $H(2^{\omega_1+})$, it is a definable class within this structure and hence $\phi_i(\vec{N}, \mathcal{F}, \mathcal{F}^\perp, \tau)$ satisfies the hypothesis of $\text{PFA}(2^{\omega_1})$. Applying Miyamoto's reformulation of $\text{PFA}(2^{\omega_1})$, there is an elementary submodel M of $H(2^{\omega_1+})$ of size ω_1 which contains ω_1 as a subset, \mathcal{F} , \mathcal{F}^\perp , τ as elements and is such that $H(\omega_2)$ satisfies there is an \vec{N} such that $\phi_i(\vec{N}, \pi(\mathcal{F}), \pi(\mathcal{F}^\perp), \pi(\tau))$. Here π is the transitive collapse of M . Notice that

since ω_1 is a subset of M , the collapsing map fixes elements of $H(\omega_2)$. In particular, it fixes τ and elements of \mathcal{F} and \mathcal{F}^\perp . It follows that the postulated \vec{N} really is a reflecting sequence for $\Sigma^i_{\mathcal{F}}$.

Now fix, for each i , a reflecting sequence $\langle M_\nu^i : \nu < \omega_1 \rangle$ for $\Sigma^i_{\mathcal{F}}$. Let E be the collection of all ν such that $M_\nu^i \cap \omega_1 = \nu$, for each i . Letting

$$N_\nu = (\mathcal{F} \cup \mathcal{F}^\perp) \cap \bigcup_{i < \omega} M_\nu^i,$$

it is easily checked that $\langle N_\nu : \nu \in E \rangle$ is a witness to $\varphi(\mathcal{F})$. ■

LEMMA 3.8: *For a given family \mathcal{F} of subtrees of an Aronszajn tree T , there is a proper forcing extension which satisfies $\varphi(\mathcal{F})$.*

Proof. Construct a countable support iteration of length ω such that at the i -th stage of the iteration, a reflecting sequence is added to $\dot{\Sigma}^i_{\mathcal{F}}$ by a proper forcing. It is easily checked that the iteration generates a generic extension which satisfies $\varphi(\mathcal{F})$. ■

Remark 3.9: The reader is cautioned that it does not immediately follow that if \mathcal{F} is moreover predense, then there is a proper forcing extension in which $\psi(\mathcal{F})$ holds, since a priori \mathcal{F} may fail to be predense in the generic extension. This is addressed in the next section. Similarly, if \mathcal{F} is defined by a Σ_1 -formula, then there are two versions of \mathcal{F} in a generic extension — $\hat{\mathcal{F}}$ and $\check{\mathcal{F}}$. This lemma only implies that $\phi(\hat{\mathcal{F}})$ can be forced.

COROLLARY 3.10: *If $\text{PFA}(\omega_1)$ holds and T is a saturated A-tree, then $\varphi(\mathcal{F})$ is true for all families \mathcal{F} of subtrees of T .*

Proof. Let T be an A-tree and let \mathcal{F} be a family of subtrees of \mathcal{F} . Applying Lemma 3.5, fix a subfamily \mathcal{F}' of \mathcal{F} of cardinality at most ω_1 such that $\mathcal{F}^\perp = (\mathcal{F}')^\perp$. Then $\varphi(\mathcal{F}')$ implies $\varphi(\mathcal{F})$, and $\varphi(\mathcal{F}')$ is a Σ_1 -statement in a parameter listing T and the members of \mathcal{F}' . Theorem 3.7 shows that there is a proper forcing making this Σ_1 -statement hold. ■

4. Forcing instances of ψ

As already noted, Lemma 3.8 comes short of showing that, for a given predense \mathcal{F} , there is a proper forcing extension in which $\psi(\mathcal{F})$ holds. Upon forcing a reflecting sequence for $\Sigma^0_{\mathcal{F}}$, \mathcal{F} may fail to be predense.

A similar problem arises in the context of $\mathcal{P}(\omega_1)/\text{NS}$. For a given antichain \mathcal{A} in $\mathcal{P}(\omega_1)/\text{NS}$, there is a stationary subset S of $[\omega_1 \cup \mathcal{A}]^\omega$ such that $\psi_{\text{NS}}(\mathcal{A})$ is equivalent to S strongly reflecting in the sense of [5, p. 57]. Furthermore, there is a semi-proper forcing \mathcal{Q} such that if generic absoluteness holds for \mathcal{Q} in the sense of the previous section, then S strongly reflects. This does not ensure, however, that $\psi_{\text{NS}}(\mathcal{A})$ holds after forcing with \mathcal{Q} . In fact, while semi-proper forcing can always be iterated with revised countable support while preserving ω_1 , there are models such as L in which there is no set forcing which makes $\mathcal{P}(\omega_1)/\text{NS}$ saturated. Hence, in the case of $\mathcal{P}(\omega_1)/\text{NS}$, the discrepancy between forcability and the consequences of generic absoluteness can represent an insurmountable difficulty.

In this section we will see that the saturation of A-trees is fundamentally different in this regard. We will show that there is a single set mapping associated with a given predense \mathcal{F} such that if the set mapping reflects, $\psi(\mathcal{F})$ is true.

LEMMA 4.1: *Suppose that $n \in \omega$, T is an A-tree, \mathcal{F} is a predense collection of subtrees of T and M is a countable elementary submodel of $H((\beth_{n+2})^+)$ which contains T and \mathcal{F} as elements. Let $\delta = M \cap \omega_1$. Suppose X is an n -element subset of the δ -th level. Then*

$$\{P \in H(\omega_2) : \forall t \in X \exists A \in \mathcal{F} \cap P (t \upharpoonright (P \cap \omega_1) \in A)\}$$

is M -stationary.

Proof. We prove this by an induction on n . In the base case $n = 0$, there is nothing to show. Now suppose that the lemma is true for n and let M be a countable elementary submodel of $H((\beth_{n+3})^+)$ and X be an $n + 1$ -element subset of the δ -th level of T . Let E be a given closed unbounded subset of $[H(\omega_2)]^\omega$ which is in M . Let t be any element of X and let $X_0 = X \setminus \{t\}$. The set of elements of $[H(\omega_2)]^\omega$ of the form $N \cap H(\omega_2)$, for some countable elementary submodel N of $H((\beth_{n+2})^+)$, is a club set in M , so, applying Lemma 3.6, there is a countable elementary submodel N of $H((\beth_{n+2})^+)$ such that \mathcal{F}, E are in N and there is an A in $\mathcal{F} \cap N$ such that $t \upharpoonright (N \cap \omega_1)$ is in A . Let E^* be the set of all P in E such that A is in P . Clearly, E^* is a club and belongs to N . Applying the inductive hypothesis to N and X_0 , there is a P in $E^* \cap N$ such that for every s in X_0 , there is a A' in $\mathcal{F} \cap P$ such that $s \upharpoonright (P \cap \omega_1)$ is in A' . But A is also in P and $t \upharpoonright (P \cap \omega_1)$ is in A as well since it is downward closed. ■

LEMMA 4.2: *If \mathcal{F} is a predense family of subtrees of T , then there is an open stationary set mapping $\Sigma_{\mathcal{F}}$ such that if $\Sigma_{\mathcal{F}}$ admits a reflecting sequence, then $\psi(\mathcal{F})$ is true.*

Proof. Fix, for each limit $\alpha < \omega_1$, a cofinal $C_\alpha \subseteq \alpha$ of order type ω . Let $T = \{t(\alpha, i) : \alpha < \omega_1 \text{ and } i < \omega\}$ be such that for every α and i , the height of $t(\alpha, i)$ is α . If M is a countable elementary submodel of $H((\beth_{\omega+1})^+)$, define $\Sigma_{\mathcal{F}}(M)$ as follows. Let $\delta = M \cap \omega_1$ and let $\Sigma_{\mathcal{F}}(M)$ be the set of all $P \in [H(\omega_2)]^\omega$ such that either P is not an elementary submodel of $H(\omega_2)$ or else for every $i < |C_{M \cap \omega_1} \cap P|$ there is an A in $P \cap \mathcal{F}$ which contains $t(\delta, i) \upharpoonright (P \cap \omega_1)$.

It is easily checked that $\Sigma_{\mathcal{F}}(M)$ is open for every M . It should be clear that a reflecting sequence of $\Sigma_{\mathcal{F}}$ can easily be modified to produce a witness $\langle N_\nu : \nu \in E \rangle$ to $\psi(\mathcal{F})$. Therefore, it remains to show that $\Sigma_{\mathcal{F}}(M)$ is M -stationary for all M in the domain of $\Sigma_{\mathcal{F}}$. To see this, let $E \subseteq H(\omega_2)$ be a club. Find a countable elementary submodel N of $H((\beth_\omega)^+)$ which is an element of M and contains E as a member. Denote $n = |C_{M \cap \omega_1} \cap N|$ and apply Lemma 4.1 to N and n to find a P in $E \cap \Sigma_{\mathcal{F}}(M) \cap M$. ■

LEMMA 4.3: *If \mathcal{F} is a predense family of subtrees of an A -tree, then there is a proper forcing extension in which $\psi(\mathcal{F})$ holds.*

Proof. By the proof of Theorem 3.1 of [12], there is a proper forcing which adds a reflecting sequence to the $\Sigma_{\mathcal{F}}$ of Lemma 4.2. ■

5. Relative consistency results

In this section we will present a number of iterated forcing constructions aimed at proving upper bounds on the consistency of φ and the existence of a five element basis for the uncountable linear orders. Throughout this section we will utilize the following standard facts about L .

THEOREM 5.1 (see [6, 2.2]): *Suppose $V = L$. If κ is an uncountable regular cardinal and E is a stationary subset of κ , then $\diamond_E(\kappa)$ holds: there is a sequence $\langle A_\xi : \xi \in E \rangle$ such that for all $X \subseteq \kappa$,*

$$\{\xi \in E : X \cap \xi = A_\xi\}$$

is stationary.

Remark 5.2: If $\diamond_E(\kappa)$ holds and every element of E is an inaccessible cardinal, then $\diamond_E(\kappa)$ is equivalent to the following stronger statement: There is a sequence $\langle A_\delta : \delta \in E \rangle$ of elements of $H(\kappa)$ such that if X_i ($i < n$) is a finite sequence of subsets of $H(\kappa)$, then there is a stationary set of δ in E such that

$$A_\delta = \langle X_i \cap H(\delta) : i < n \rangle.$$

It is easily checked that if κ is a regular cardinal in V , then it is also a regular cardinal in L . Hence, if κ is inaccessible (Mahlo), then L satisfies that κ is inaccessible (Mahlo). Reflecting cardinals also relativize to L [8].

THEOREM 5.3: *Suppose that there is a Mahlo cardinal. Then there is a forcing extension of L which satisfies φ .*

Proof. Let κ be Mahlo and note that κ is also Mahlo in L ; from now on, work in L . Let E be the stationary set of inaccessible cardinals less than κ and, applying Theorem 5.1, let $\langle A_\delta : \delta \in E \rangle$ be a $\diamond_E(\kappa)$ -sequence in the revised sense stated in Remark 5.2. Construct a countable support iteration $\langle \mathcal{P}_\alpha; \mathcal{Q}_\alpha : \alpha < \kappa \rangle$ of proper forcing notions of size $< \kappa$. If $\alpha \in E$ and $A_\alpha = \langle \dot{T}, \dot{\mathcal{F}} \rangle$, where \dot{T} is a \mathcal{P}_α -name for an A-tree and $\dot{\mathcal{F}}$ is a \mathcal{P}_α -name for a family of subtrees of \dot{T} , then we let $\dot{\mathcal{Q}}_\alpha$ be a proper forcing in $H(\kappa)$ which first forces $\psi(\dot{\mathcal{F}} \cup \dot{\mathcal{F}}^\perp)$ and then forces $\varphi(\dot{\mathcal{F}})$. In other cases we can let $\dot{\mathcal{Q}}_\alpha$ be any proper forcing in $H(\kappa)$. Let \mathcal{P}_κ be the limit of the iteration. By standard arguments the forcing \mathcal{P}_κ is proper and κ -c.c. [9].

Suppose now \dot{T} is a \mathcal{P}_κ -name for an A-tree and $\dot{\mathcal{F}}$ is a \mathcal{P}_κ -name for a family of subtrees of \dot{T} . Let $\dot{\mathcal{F}}_\delta$ be the set of all \mathcal{P}_δ -names \dot{S} which are forced by every condition to be in $\dot{\mathcal{F}}$. Since κ is Mahlo and each of the iterands of \mathcal{P}_κ has cardinality less than κ , there is a relative closed and unbounded set D of δ in E , such that \mathcal{P}_δ has the δ -c.c., \dot{T} is a \mathcal{P}_δ -name, and if \dot{S} is a \mathcal{P}_δ -name for a subtree of \dot{T} which has countable intersection with every element of $\dot{\mathcal{F}}_\delta$, then \dot{S} is forced to be in $\dot{\mathcal{F}}^\perp$. Since $\langle A_\alpha : \alpha \in E \rangle$ is a $\diamond_E(\kappa)$ -sequence, there is a δ in D such that $A_\delta = \langle \dot{T}, \dot{\mathcal{F}}_\delta \rangle$. At stage δ the partial order $\dot{\mathcal{Q}}_\delta$ forces both $\psi(\dot{\mathcal{F}}_\delta \cup \dot{\mathcal{B}})$ and $\varphi(\dot{\mathcal{F}}_\delta)$, where $\dot{\mathcal{B}}$ is $\dot{\mathcal{F}}_\delta^\perp$ computed after forcing with \mathcal{P}_δ . By choice of δ , $\dot{\mathcal{B}}$ is forced to be a subset of $\dot{\mathcal{F}}^\perp$. Since $\psi(\dot{\mathcal{F}}_\delta \cup \dot{\mathcal{B}})$ is a Σ_1 -formula, it is upwards absolute and hence forced by \mathcal{P}_κ . By Lemma 2.2, \mathcal{P}_κ forces $\dot{\mathcal{F}}_\delta \cup \dot{\mathcal{B}}$ is predense and hence that $\dot{\mathcal{F}}_\delta^\perp = \dot{\mathcal{F}}^\perp$. Consequently, $\varphi(\dot{\mathcal{F}}_\delta)$ implies $\varphi(\dot{\mathcal{F}})$. ■

THEOREM 5.4: *If there is a cardinal which is both reflecting and Mahlo, then there is a proper forcing extension of L which satisfies the conjunction of $\text{PFA}(\omega_1)$ and φ . In particular, the forcing extension satisfies that the uncountable linear orders have a five element basis.*

Proof. This is very similar to the proof of Theorem 5.3, except that at stages $\alpha < \kappa$ which are not in E , we force with partial orders in $H(\kappa)$ given by an appropriate book keeping device. Following [8], it is possible to arrange that $\text{PFA}(\omega_1)$ holds in the generic extension as well. ■

THEOREM 5.5: *Suppose that there is an inaccessible cardinal κ such that for every $\kappa_0 < \kappa$, there is an inaccessible cardinal $\delta < \kappa$ such that κ_0 is in $H(\delta)$ and $H(\delta)$ satisfies that there are two reflecting cardinals which are greater than κ_0 . Then there is a proper forcing extension in which κ is ω_2 and the uncountable linear orders have a five element basis.*

First observe that by taking a direct sum of trees, Theorems 4.2 implies that if \vec{T} is an ω -sequence of A-trees and $\vec{\mathcal{F}}$ is an ω -sequence such that \mathcal{F}_n is a family of subtrees of T_n , then there is a single set mapping $\Sigma_{\vec{\mathcal{F}}}$ such that if $\Sigma_{\vec{\mathcal{F}}}$ admits a reflecting sequence, then $\psi(\mathcal{F}_n \cup \mathcal{F}_n^\perp)$ holds for all $n < \omega$. Theorem 5.5 can be proved by iterating the forcings provided by the following lemmas with appropriate book keeping. By mixing in appropriate σ -closed collapsing forcings as needed, we may ensure that the iteration has the κ -c.c. but collapses every uncountable cardinal less than κ to ω_1 .

LEMMA 5.6 (see [18, §4.4]): *Let L be an uncountable linear order, X be a set of reals of size \aleph_1 , and C be a Countryman order. There is a proper forcing \mathcal{P} of cardinality less than \beth_ω which forces that “ L contains an isomorphic copy of X , ω_1 , ω_1^* , C , or C^* ” is equivalent to an instance of CTA.*

LEMMA 5.7: *Suppose that T is an A-tree, K is a subset of T , and there is an inaccessible δ such that $H(\delta)$ satisfies that there are two reflecting cardinals. Then there is a proper forcing in $H(\delta)$ which forces the instance of CTA for K and T .*

Proof. If λ is a reflecting cardinal in $H(\delta)$, let \mathcal{P}_λ denote the proper forcing which satisfies the λ -c.c. and forces that $H(\delta)^{V^{\mathcal{P}_\lambda}}$ satisfies $\text{PFA}(\omega_1)$. If $\vec{\mathcal{F}}$ is an ω -sequence of families of subtrees of A-trees, let $\mathcal{Q}_{\vec{\mathcal{F}}}$ be the proper forcing which forces the conjunction of $\psi(\mathcal{F}_n \cup \mathcal{F}_n^\perp)$ and $\varphi(\mathcal{F}_n)$ for all n . Let $\lambda_0 < \lambda_1$

be the two reflecting cardinals in $H(\delta)$. We claim that

$$(\mathcal{P}_{\lambda_0} * \dot{\mathcal{Q}}_{\mathcal{R}}) * \dot{\mathcal{P}}_{\lambda_1}$$

is the desired proper forcing, where \mathcal{R}_n is the family of subtrees of $T^{[n]}$ defined in Section 2. Clearly this forcing is proper and an element of $H(\delta)$. It suffices to show that it forces the instance of CTA for T and K . The key observation is that, after forcing with \mathcal{P}_{λ_0} , if S is an element of \mathcal{R}_n^\perp for some n , then S remains in \mathcal{R}_n^\perp after any proper forcing which is in $H(\delta)^{V^{\mathcal{P}_{\lambda_0}}}$. This is because asserting that S is not in \mathcal{R}_n^\perp is a Σ_1 -statement with parameters T , K , and S . By arguments given in the proof of Theorem 5.3, $\mathcal{Q}_{\mathcal{R}}$ forces $\varphi(\mathcal{R}_n)$ to be true for all n and, moreover, that this statement remains true after further forcing with \mathcal{P}_{λ_1} . Applying Lemma 2.7 in the extension by

$$(\mathcal{P}_{\lambda_0} * \dot{\mathcal{Q}}_{\mathcal{R}}) * \dot{\mathcal{P}}_{\lambda_1},$$

both $\text{PFA}(\omega_1)$ and the Key Lemma for T and K hold. Therefore, by theorems from [13], the instance of CTA for T and K is true. ■

6. Concluding remarks and questions

Observe that the property of κ in the statement of Theorem 5.5 is expressible by a Σ_0 -formula with no parameters. Hence the least such cardinal is not reflecting and it is, therefore, possible, if such cardinals exist at all, to produce a forcing extension of L in which Shelah’s conjecture is true and ω_2 is not reflecting in L . On the other hand, we do not know the answer to the following.

QUESTION 6.1: *Suppose that the uncountable linear orders have a five element basis. Is there a $\delta < \omega_2$ such that L_δ satisfies “there is a reflecting cardinal?”*

It is also natural to ask:

QUESTION 6.2: *Does $\text{PFA}(\omega_1)$ imply Aronszajn tree saturation?*

The only known direct construction of a failure of A-tree saturation is given in [4, §2] and is based upon the existence of a Kurepa tree. Baumgartner has shown that $\text{PFA}(\omega_1)$ implies that there are no Kurepa trees [2, 7.11].⁷ This is likely closely related to the consistency strength of φ .

⁷ The hypothesis which appears in [2, 7.11] is PFA but the proof shows that the conclusion follows from $\text{PFA}(\omega_1)$.

QUESTION 6.3: *If φ is true, must ω_2 be Mahlo in L ?*

References

- [1] U. Abraham and S. Shelah, *Isomorphism types of Aronszajn trees*, Israel Journal of Mathematics, **50** (1985), 75–113.
- [2] J. E. Baumgartner, *Applications of the Proper Forcing Axiom*, in *Handbook of Set-Theoretic Topology*, (K. Kunen and J. Vaughan, eds.), North-Holland, 1984.
- [3] J. E. Baumgartner, *A new class of order types*, Annals of Mathematical Logic **9** (1976), 187–222.
- [4] J. E. Baumgartner, *Bases for Aronszajn trees*, Tsukuba Journal of Mathematics, **9** (1985), 31–40.
- [5] M. Bekkali, *Topics in Set Theory*. Springer-Verlag, Berlin, 1991.
- [6] Keith J. Devlin, *Constructibility*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1984.
- [7] M. Foreman, M. Magidor and S. Shelah, *Martin’s Maximum, saturated ideals, and non-regular ultrafilters. I*, Annals of Mathematics (2) **127** (1988), 1–47.
- [8] M. Goldstern and S. Shelah, *The Bounded Proper Forcing Axiom*, Journal of Symbolic Logic **60**(1995), 58–73.
- [9] T. Jech, *Set Theory*, Perspectives in Mathematical Logic, second edition, Springer-Verlag, Berlin, 1997.
- [10] T. Miyamoto, *A note on weak segments of PFA*, in *Proceedings of the Sixth Asian Logic Conference* (Beijing, 1996), World Sci. Publishing, 1998, pp. 175–197.
- [11] J. T. Moore, *Structural analysis of Aronszajn trees*, Proceedings of the 2005 Logic Colloquium, Athens, Greece.
- [12] J. T. Moore, *Set mapping reflection*, Journal of Mathematical Logic **5** (2005), 87–97.
- [13] J. T. Moore, *A five element basis for the uncountable linear orders*, Annals of Mathematics (2) **163** (2006), 669–688.
- [14] S. Shelah, *Decomposing uncountable squares to countably many chains*, Journal Combinatorial Theory Ser. A **21** (1976), 110–114.
- [15] S. Todorćević, *Lipschitz maps on trees*, report 2000/01 number 13, Institut Mittag-Leffler.
- [16] S. Todorćević, *Trees and linearly ordered sets*, in *Handbook of set-theoretic topology*, North-Holland, Amsterdam, 1984, pp. 235–293.
- [17] S. Todorćević, *Localized reflection and fragments of PFA*, in *Logic and scientific methods*, volume 259 of DIMACS Ser., Discrete Math. Theoret. Comput. Sci., AMS, 1997, pp. 145–155.
- [18] S. Todorćević, *Walks on Ordinals and their Characteristics*, Progress in Mathematics, 263, Birkhäuser, Basel, 2007.